arXiv:0706.0790v2 [math.AT] 3 Mar 2009

FREE ACTIONS OF FINITE GROUPS ON $S^n \times S^n$

IAN HAMBLETON AND ÖZGÜN ÜNLÜ

Abstract. Let p be an odd prime. We construct a non-abelian extension Γ of S^1 by $\mathbf{Z}/p \times \mathbf{Z}/p$, and prove that any finite subgroup of Γ acts freely and smoothly on $S^{2p-1} \times S^{2p-1}$. In particular, for each odd prime p we obtain free smooth actions of infinitely many non-metacyclic rank two p-groups on $S^{2p-1} \times S^{2p-1}$. These results arise from a general approach to the existence problem for finite group actions on products of equidimensional spheres.

Introduction

Conner [14] and Heller [18] proved that any finite group G acting freely on a product of two spheres must have $\operatorname{rank} G \leqq 2$. In other words, the maximal rank of an elementary abelian subgroup of G is at most two. However, if both spheres have the same dimension then there are additional restrictions: the alternating group A_4 of order 12 has rank two, but does not admit such an action (see Oliver [31]). It was observed by Adem-Smith [3, p. 423] that A_4 is a subgroup of every rank two simple group, so all these are ruled out.

Question. *What group theoretic conditions characterize the rank two finite groups which can act freely and smoothly on $S^n \times S^n$, for some $n \geqq 1$?*

The work of G. Lewis [27] shows that for every prime p, the p-Sylow subgroup of a finite group G which acts freely on $S^n \times S^n$ is abelian unless $n = 2pr - 1$ for some $r \geq 1$. Lewis also points out [27, p. 538] that metacyclic groups act freely and smoothly on some $S^n \times S^n$, but the existence of a free action by any other non-abelian p-group, for p odd, has been a long-standing open question. In this paper we provide a general approach to this problem, and construct an infinite family of new examples for each odd prime in the minimal dimension.

For each odd prime p, let Γ be the Lie group given by the following presentation

$$\Gamma = \left\langle a, b, z \mid z \in S^1,\ a^p = b^p = [a, z] = [b, z] = 1,\ [a, b] = \omega \right\rangle$$

where $\omega = e^{2\pi i/p} \in S^1 \subseteq \mathbb{C}$. This is a non-abelian central extension of $\mathbf{Z}/p \times \mathbf{Z}/p$ by S^1.

Theorem A. *Let p be an odd prime, and let G be a finite subgroup of Γ. Then G acts freely and smoothly on $S^{2p-1} \times S^{2p-1}$.*

Date: Jan. 14, 2009.
Research partially supported by NSERC Discovery Grant A4000.

The finite subgroups of Γ which surject onto the quotient $\mathbf{Z}/p\times\mathbf{Z}/p$ are direct products $G = C \times P(k)$, where C is a finite cyclic group of order prime to p, and

$$P(k) = \Big\langle a, b, c \,|\, a^p = b^p = c^{p^{k-2}} = [a, c] = [b, c] = 1,\ [a, b] = c^{p^{k-3}} \Big\rangle$$

is a rank two p-group of order p^k, $k \geq 3$. We therefore obtain infinitely many actions of non-metacyclic p-groups on $S^{2p-1} \times S^{2p-1}$ for each prime p.

An important special case is the extraspecial p-group $G_p = P(3)$ of order p^3 and exponent p. Our existence result contradicts claims made in [4], [5], [37], and [41] that G_p-actions do not exist (for cohomological reasons) on any product of equidimensional spheres. It was later shown by Benson and Carlson [7] that such actions could not be ruled out for any prime p by cohomological methods. Moreover for $p = 3$, in [17], we gave an explicit construction of a free smooth action of Γ (and in particular G_3) on $S^5 \times S^5$. This construction provides an alternate proof of Theorem A for $p = 3$.

More generally, rank two finite p-groups were classified by Blackburn [8] (see also [26]). Consider the additional family, extending the groups $P(k)$:

$$B(k, \epsilon) = \Big\langle a, b, c \,|\, a^p = b^p = c^{p^{k-2}} = [b, c] = 1,\ [a, c] = b,\ [a, b] = c^{\epsilon p^{k-3}} \Big\rangle$$

where $k \geq 4$, and ϵ is 1 or a quadratic non-residue $\bmod p$. Here is Blackburn's list of the rank two p-groups G with order p^k, and $p > 3$ (the classification for $p = 3$ is more complicated):

I) G is a metacyclic p-group.
II) $G = P(k)$, for $k \geq 3$.
III) $G = B(k, \epsilon)$, for $k \geq 4$.

We now know that groups of types I and II do act freely on a product of equidimensional spheres in the minimal dimension. Is this the complete answer ?

Conjecture. *Let $p > 3$ be an odd prime. If G is a rank two p-group G which acts freely and smoothly on $S^{2pr-1} \times S^{2pr-1}$, $r \geq 1$, then G is metacyclic or G is a subgroup of Γ.*

If this conjecture is true, then we would know all the possible p-Sylow subgroups ($p > 3$) for finite groups acting freely on products of equidimensional spheres. This would be an important step forward in understanding the general problem.

We remark that in order to handle groups of composite order, it is necessary to establish the existence of free actions of p-groups in higher dimensions $S^{2pr-1} \times S^{2pr-1}$, $r > 1$. In [17], we discussed this existence problem specifically for $p = 3$, $r = 2$, and showed that all odd order subgroups of $SU(3)$, including the extraspecial 3-group G_3 and the type III group $B(4, -1)$, can act freely and smoothly on $S^{11} \times S^{11}$. In particular, we are suggesting that existence results for $p = 3$ will be qualitatively different than those for $p > 3$.

We can expect an even more complicated structure for the 2-Sylow subgroup of a finite group acting freely on some $S^n \times S^n$, since this is already the case for free actions on S^n. We can take products of periodic groups $G_1 \times G_2$ and obtain a variety of actions of non-metacyclic groups on $S^n \times S^n$ (see [16] for the existence of these examples, generalizing the results of Stein [34]). Here the 2-groups are all metabelian, so one might hope that this is the correct restriction on the 2-Sylow subgroup. However, there are non-metabelian 2-groups which are subgroups of $Sp(2)$, hence by generalizing the notion of fixity in [2] to quaternionic fixity, one can construct free actions of these non-metabelian 2-groups on $S^7 \times S^7$ (see [38]).

Remark. Every rank two finite p-group (for p odd) admits a free smooth action on some product $S^n \times S^m$, $m \geqq n$ (see [2] for $p > 3$, [38] for $p = 3$). The survey article by A. Adem [1] describes recent progress on the existence problem in this setting for general finite groups (see also [20]). In most cases, construction of the actions requires $m > n$.

We will always assume that our actions on $S^n \times S^n$ are homologically trivial and n is odd. For free actions of odd p-groups this follows from the Lefschetz fixed point theorem.

Acknowledgement. The authors would like to thank Alejandro Adem, Dave Benson, Jim Davis and Matthias Kreck for useful conversations and correspondence.

1. An overview of the proof

Given a group G, and two cohomology classes θ_1, $\theta_2 \in H^{n+1}(G; \mathbf{Z})$, we can construct an associated space B_G as the total space of the induced fibration

$$\begin{tikzcd}
K(\mathbf{Z} \oplus \mathbf{Z}, n) \arrow[r] & B_G \arrow[d] & \\
 & BG \arrow[r, "\theta_1, \theta_2"] & K(\mathbf{Z} \oplus \mathbf{Z}, n+1)
\end{tikzcd}$$

where BG denotes the classifying space of G. If we also have a stable oriented bundle $\nu_G \colon B_G \to BSO$, then we can consider the bordism groups

$$\Omega_k(B_G, \nu_G)$$

defined as in [35, Chap. II]. The objects are commutative diagrams

$$\begin{tikzcd}
\nu_M \arrow[r, "b"] \arrow[d] & \nu_G \arrow[d] \\
M \arrow[r, "f"] & B_G
\end{tikzcd}$$

where M^k is a closed, smooth k-dimensional manifold with stable normal bundle ν_M, $f \colon M \to B_G$ is a reference map, and $b \colon \nu_M \to \nu_G$ is a stable bundle map covering f. The bordism relation is the obvious one consistent with the normal data and reference maps.

Our general strategy is to use the space B_G as a model for the n-type of the orbit space of a possible free G-action on $S^n \times S^n$, and ν_G as a candidate for its stable normal bundle. For G finite, the actual orbit space (a closed $2n$-dimensional manifold) is obtained by surgery on a representative of a suitable bordism element in $\Omega_{2n}(B_G, \nu_G)$. This approach to the problem follows the general outline of Kreck's "modified surgery" program (see [23]).

We will carry out this strategy uniformly to prove Theorem A. We may restrict our attention to the finite subgroups $G \subset \Gamma$ which surject onto $\mathbf{Z}/p \times \mathbf{Z}/p$. Then, by construction, the induced map on classifying spaces gives a circle bundle

$$S^1 \to B_G \to B_\Gamma \ .$$

Let $n = 2p-1$. We select appropriate data $(\theta_1, \theta_2, \nu_\Gamma)$ for Γ, and then define the data for each $G \subset \Gamma$ by restriction. We study B_Γ and the bordism groups $\Omega_{2n-1}(B_\Gamma, \nu_\Gamma)$ to carry out the following steps.

(i) We construct a non-empty subset $T_\Gamma \subseteq H_{2n-1}(B_\Gamma; \mathbf{Z})$, depending on the data $(\theta_1, \theta_2, \nu_\Gamma)$, consisting entirely of primitive elements of infinite order.

(ii) For each $\gamma \in T_\Gamma$, we show that there is a bordism element $[N, c] \in \Omega_{2n-1}(B_\Gamma, \nu_\Gamma)$ whose image $c_*[N] = \gamma \in T_\Gamma$ under the Hurewicz map $\Omega_{2n-1}(B_\Gamma, \nu_\Gamma) \to H_{2n}(B_\Gamma; \mathbf{Z})$.

One of the key points is that the cohomology of the groups Γ is much simpler than that of its finite subgroups (see Leary [24] for Γ, and Lewis [28] for the extra-special p-groups), so the computations of Steps (i) and (ii) are best done over Γ.

Now for each finite subgroup $G \subset \Gamma$ as above, define T_G as the image of T_Γ under the S^1-bundle transfer

$$trf\colon H_{2n-1}(B_\Gamma; \mathbf{Z}) \to H_{2n}(B_G; \mathbf{Z})$$

induced by the fibration of classifying spaces. The subset T_G will contain the images of fundamental classes of the possible free G-actions on $S^n \times S^n$. For each $\gamma_G = trf(\gamma) \in T_G$, we have a bordism element $[M, f] \in \Omega_{2n}(B_G, \nu_G)$, where M is the total space of the pulled-back S^1-bundle over $[N, c]$ with $c_*[N] = \gamma$. We then show that we can obtain $\widetilde{M} = S^n \times S^n$ by surgery on $[M, f]$ within its bordism class.

2. REPRESENTATIONS AND COHOMOLOGY OF Γ

§2A. Some subgroups of Γ. For each odd prime p, the Lie group Γ is a central extension

$$1 \to S^1 \to \Gamma \to Q_p \to 1$$

where $Q_p = \mathbf{Z}/p \times \mathbf{Z}/p$. We fix the presentation for Γ given in the Introduction, with generators $\langle a, b\rangle$ for Q_p. For any finite subgroup $G \subset \Gamma$ which surjects onto Q_p, we have a commutative diagram of central extensions

$$\begin{tikzcd}
1 \arrow[r] & \langle c\rangle \arrow[r] \arrow[d] & G \arrow[r] \arrow[d] & Q_p \arrow[r] \arrow[d, equal] & 1 \\
1 \arrow[r] & S^1 \arrow[r] & \Gamma \arrow[r] & Q_p \arrow[r] & 1
\end{tikzcd}$$

where the centre $Z(G) = \langle c\rangle \subset G$ is a finite cyclic group. Now we list some subgroups of Γ which will be important in our calculations.

Definition 2.1. Let $d_t = ab^t$ if $0 \le t \le p-1$, $d_p = b$, and define $D_t = \langle d_t\rangle$, for $0 \le t \le p$. Let $\Sigma_t = \langle d_t, S^1\rangle$ denote the subgroup of Γ generated by d_t and S^1 for $0 \le t \le p$.

We will usually write Σ instead of Σ_p for the subgroup of Γ generated by b and S^1.

Remark 2.2. Since the subgroup $S^1 \subset \Gamma$ is central, any continuous group automorphism $\phi \in \mathrm{Aut}(\Gamma)$ induces an automorphism $\bar{\phi} \in \mathrm{Aut}(Q_p) = GL_2(p)$. In Section 6, we will use the fact that the image of $\mathrm{Aut}(\Gamma)$ contains the subgroup $SL_2(p) \subset GL_2(p)$. More explicitly, for each matrix $A \in SL_2(p)$, we can define an automorphism $\phi_A \in \mathrm{Aut}(\Gamma)$ such that $\bar{\phi}_A = A$ as follows. Any element in Γ can be written as $a^r b^s z$ for unique $r, s \in \mathbf{Z}/p$ and $z \in S^1$. Given a matrix $A = \begin{pmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \end{pmatrix}$ in $SL_2(p)$ we can define $\phi_A(a) = a^{r_{11}} b^{r_{12}}$, $\phi_A(b) = a^{r_{21}} b^{r_{22}}$ and $\phi_A(z) = z$. By construction, $\bar{\phi}_A = A$.

§2B. Representations of Γ and some of its subgroups. First, we define a 1–dimensional representation

$$\Phi_t \colon \Gamma \to U(1), \quad \text{for } 0 \le t \le p,$$

so that $\ker \Phi_t = \Sigma_{p-t}$ and $\Phi_t(d_t) = e^{2\pi i/p}$. For any subgroup $G \subset \Gamma$ and $0 \le t \le p$ we define a 1-dimensional representation of G by the formula:

$$\Phi_{t,G} = \mathrm{Res}^{\Gamma}_{G}(\Phi_t) : G \to U(1), \quad 0 \le t \le p \ .$$

Second, we define a 1-dimensional representation Φ'_t of Σ_t by setting:

$$\Phi'_t \colon \Sigma_t \to U(1), \quad 0 \le t \le p,$$

where $\Phi'_t(d_t) = 1$ and $\Phi'_t(z) = z$ for z in S^1. For any subgroup $G \subset \Sigma_t$ we define a 1-dimensional representation of G by the formula:

$$\Phi'_{t,G} = \mathrm{Res}^{\Sigma_t}_{G}(\Phi'_t) : G \to U(1), \quad 0 \le t \le p \ .$$

Finally, we define a p–dimensional irreducible representation Ψ of Γ as follows:

$$\Psi = \mathrm{Ind}^{\Gamma}_{\Sigma}(\Phi'_p) \colon \Gamma \to SU(p)$$

and for any subgroup $G \subset \Gamma$ we define:

$$\Psi_G = \mathrm{Res}^{\Gamma}_{G}(\Psi) : G \to SU(p)$$

by restriction, as a p-dimensional representation of G.

§2C. Cohomology of Γ and some of its subgroups. We will use the notations and results of Leary [24] for the integral cohomology ring of Γ.

Theorem 2.3 ([24, Theorem 2]). *$H^*(B\Gamma; \mathbf{Z})$ is generated by elements $\alpha, \beta, \sigma_1, \chi_2, \dots, \chi_{p-1}, \zeta$, with*

$$\deg(\alpha) = \deg(\beta) = 2, \quad \deg(\zeta) = 2p, \quad \deg(\chi_i) = 2i,$$

subject to some relations.

In the statement of Theorem 2.3, the elements $\alpha = \Phi_0 \colon \Gamma \to U(1)$ and $\beta = \Phi_p \colon \Gamma \to U(1)$, by considering $H^2(B\Gamma; \mathbf{Z}) = \mathrm{Hom}(\Gamma, S^1)$, and ζ is the p^{th} Chern class of the p–dimensional irreducible representation Ψ of Γ. The mod p cohomology ring of Γ is also given by Leary:

Theorem 2.4 ([25, Theorem 2]). *$H^*(B\Gamma;\mathbf{Z}/p)$ is generated by elements y, y', x, x', $c_2, c_3, \dots, c_{p-1}, z$, with*

$$\begin{aligned} \deg(y) &= \deg(y') = 1,\ \deg(x) = \deg(x') = 2, \\ \deg(z) &= 2p, \quad \textit{and} \quad \deg(c_i) = 2i, \end{aligned}$$

subject to some relations.

Let π_* stand for the projection map from $H^*(B\Gamma;\mathbf{Z})$ to $H^*(B\Gamma;\mathbf{Z}/p)$, and δ_p for the Bockstein from $H^*(B\Gamma;\mathbf{Z}/p)$ to $H^{*+1}(B\Gamma;\mathbf{Z})$ then $\delta_p(y) = \alpha$, $\delta_p(y') = \beta$, $\pi_*(\alpha) = x$, $\pi_*(\beta) = x'$, $\pi_*(\chi_i) = c_i$, and $\pi_*(\zeta) = z$. Here are some facts about the cohomology of certain subgroups.

Remark 2.5. Considering $H^2(BG,\mathbf{Z}) = \mathrm{Hom}(G, S^1)$

1. $H^*(BS^1;\mathbf{Z}) = \mathbf{Z}[\tau]$ where $\tau = \Phi'_{t,S^1}$. So τ is the identity map on S^1.
2. $H^*(B\Sigma_t;\mathbf{Z}) = \mathbf{Z}[\tau', v' \,|\, pv' = 0]$ where $\tau' = \Phi'_t$ and $v' = \Phi_{t,\Sigma_t}$
3. $H^*(B\Sigma_t,\mathbf{Z}/p) = \mathbf{F}_p[\bar\tau] \otimes (\Lambda(u) \otimes \mathbf{F}_p[v])$ where $\bar\tau$ and v are mod p reductions of τ' and v' respectively and $\beta(u) = v$.

We calculate some restriction maps:

$$\mathrm{Res}^{\Gamma}_{\Sigma_t}(\alpha) = \begin{cases} v' & \text{if } \ 0 \le t \le p-1, \\ 0 & \text{if } \ t = p\ . \end{cases} \quad \text{and} \quad \mathrm{Res}^{\Gamma}_{\Sigma_t}(\beta) = \begin{cases} tv' & \text{if } \ 0 \le t \le p-1, \\ v' & \text{if } \ t = p\ . \end{cases}$$

The property

$$\mathrm{Res}^{\Gamma}_{\Sigma_t}(\alpha^p - \alpha^{p-1}\beta + \beta^p) = (v')^p,$$

for $0 \le t \le p$, shows that this element is a good candidate for a k-invariant.

3. The $(2p-1)$-type B_Γ and the bundle data

We now construct the space B_Γ needed as a model for the $(2p-1)$-type of the quotient space of our action. Then we construct a bundle ν_Γ over this space B_Γ which will pullback to the normal bundle of the quotient space of this action.

§3A. **Definition of B_Γ.** We fix the element

$$k = \theta_1 \oplus \theta_2 = \zeta \oplus \left(\alpha^p - \alpha^{p-1}\beta + \beta^p\right) \in H^{2p}(\Gamma;\mathbf{Z}) \oplus H^{2p}(\Gamma;\mathbf{Z})\ .$$

For any subgroup $G \subset \Gamma$ define

$$k_G = \mathrm{Res}^{\Gamma}_{G}(k) \in H^{2p}(G;\mathbf{Z}\oplus\mathbf{Z}),$$

and define π_G as the fibration classified by k_G:

$$\begin{tikzcd} K(\mathbf{Z}\oplus\mathbf{Z}, 2p-1) \arrow[r] & B_G \arrow[d, "\pi_G"] & \\ & BG \arrow[r, "k_G"] & K(\mathbf{Z}\oplus\mathbf{Z}, 2p)\ . \end{tikzcd}$$

Note that the natural map $BG \to B\Gamma$, induced by the inclusion, gives a diagram

$$\begin{tikzcd}
B_G \arrow[r] \arrow[d, "\pi_G"] & B_\Gamma \arrow[d, "\pi_\Gamma"] \\
BG \arrow[r] & B\Gamma
\end{tikzcd}$$

which is a pull-back square.

§3B. The bundle data over B_Γ. For any subgroup $G \subseteq \Gamma$ we will define two bundles ϖ_G and ξ_G over BG, which will pull back by the classifying map to the stable tangent and normal bundle respectively of the quotient of a possible G-action on $S^n \times S^n$. The pullbacks of these bundles over BG to bundles over B_G will be denoted by τ_G and ν_G respectively.

(1) Tangent bundles: We have the representations $\Psi_G \colon G \to SU(p)$ and $\Phi_{t,G} \colon G \to U(1)$. Let ψ_G denote the p–dimensional complex vector bundle classified by

$$\psi_G = B\Psi_G \colon BG \to BSU(p),$$

and let $\phi_{t,G}$ denote the complex line bundle classified by

$$\phi_{t,G} = B\Phi_{t,G} \colon BG \to BU(1) = BS^1 \ .$$

We define a $3p$–dimensional complex vector bundle ϖ_G on BG by the Whitney sum

$$\varpi_G = \psi_G \oplus \phi_{0,G}^{\oplus p} \oplus \phi_{p,G}^{\oplus p}$$

and use the same notation for the stable vector bundle $\varpi_G \colon BG \to BSO$. We now identify our candidate τ_G for the stable tangent bundle.

Definition 3.1. Let τ_G denote the stable vector bundle on B_G classified by the composition

$$\tau_G \colon B_G \xrightarrow{\pi_G} BG \xrightarrow{\varpi_G} BSO \ .$$

(2) Normal bundles: First we show that there is an stable inverse of the vector bundle ϖ_G over BG, when restricted to a finite skeleton of BG.

Lemma 3.2. *For any subgroup $G \subseteq \Gamma$, there exists a stable bundle $\xi_G \colon BG \to BSO$, such that $\xi_G \oplus \varpi_G = \varepsilon$, the trivial bundle, when restricted to the $(4p-1)$-skeleton of BG.*

Proof. Take $N = 4p - 1$ and let $\varpi_\Gamma|_{B\Gamma^{(N)}}$ denote the pull-back of ϖ_Γ to $B\Gamma^{(N)}$, the N–th skeleton of $B\Gamma$, by the inclusion map of $B\Gamma^{(N)}$ in $B\Gamma$. Then there exists a vector bundle ξ_Γ over $B_\Gamma^{(N)}$ such that the bundle $\xi_\Gamma \oplus (\varpi_\Gamma|_{B\Gamma^{(N)}})$ is trivial over $B\Gamma^{(N)}$, since $B\Gamma^{(N)}$ is a finite CW–complex. Stably this vector bundle is classified by a map $\xi_\Gamma \colon B\Gamma^{(N)} \to BU$ and there is no obstruction to extending this classifying map to a map $B\Gamma \to BU$, as the obstructions to doing so lie in the cohomology groups

$$H^{*+1}(B\Gamma, B\Gamma^{(N)}; \pi_*(BU)) = 0 \ .$$

We will use the same notation ξ_Γ to denote the stable vector bundle classified by any extension map $B\Gamma \to BU \to BSO$. We then define

$$\xi_G \colon BG \to BSO$$

by composition with the map $BG \to B\Gamma$ induced by $G \subseteq \Gamma$. □

We now identify our candidate ν_G for the stable normal bundle.

Definition 3.3. Let ν_G denote the stable vector bundle on B_G classified by the composition

$$\nu_G \colon B_G \xrightarrow{\pi_G} BG \xrightarrow{\xi_G} BSO \ .$$

§3C. Characteristic classes. We will now calculate some characteristic classes for the bundles ϖ_{Σ_t} and ξ_{Σ_t} over $B\Sigma_t$. The total Chern class of a bundle ξ will be denoted $c(\xi)$. See [30, p. 228] for the definition of the mod p Wu classes $q_k(\xi) \in H^{2(p-1)k}(B; \mathbf{Z}/p)$.

Lemma 3.4. *The total Chern class of* ϖ_{Σ_t} *is*

$$\mathrm{c}(\varpi_{\Sigma_t}) = \mathrm{c}(\psi_{\Sigma_t})\, \mathrm{c}(\phi_{0,\Sigma_t}^{\oplus p} \oplus \phi_{p,\Sigma_t}^{\oplus p})$$

where

1. $\mathrm{c}(\psi_{\Sigma_t}) = 1 - (v')^{p-1} + ((\tau')^p - (v')^{p-1}\tau')$
2. $\mathrm{c}(\phi_{0,\Sigma_t}^{\oplus p} \oplus \phi_{p,\Sigma_t}^{\oplus p}) = 1 + (1+t)(v')^p + t(v')^{2p}$

Proof. Given two 1-dimensional representation $\Phi \colon G \to S^1$ and $\Phi' \colon G \to S^1$ and a natural number k, we will write $\Phi^k(g) = (\Phi(g))^k$ and $(\Phi\Phi')(g) = \Phi(g)\Phi'(g)$. It is easy to see that

$$\Psi_{\Sigma_t} = \Phi'_t \oplus \Phi_{t,\Sigma_t}\Phi'_t \oplus \Phi_{t,\Sigma_t}^2\Phi'_t \oplus \cdots \oplus \Phi_{t,\Sigma_t}^{p-1}\Phi'_t \ .$$

Hence the total Chern class of ψ_{Σ_t} is

$$(1+\tau')(1+v'+\tau')(1+2v'+\tau')\dots(1+(p-1)v'+\tau') = 1-(v')^{p-1}+(\tau')^p-(v')^{p-1}\tau'$$

since $pv' = 0$.

We have $\mathrm{c}(\phi_{0,\Sigma_t}^{\oplus p}) = (1+v')^p$ when $0 \le t \le p-1$ (1 when $t = p$), and $\mathrm{c}(\phi_{p,\Sigma_t}^{\oplus p}) = (1+tv')^p$ when $0 \le t \le p-1$ (but $(1+v')^p$ when $t = p$). Hence the total Chern class of $\phi_{0,\Sigma_t}^{\oplus p} \oplus \phi_{p,\Sigma_t}^{\oplus p}$ is equal to

$$(1+v')^p(1+tv')^p = (1+(v')^p)(1+(tv')^p) = (1+(1+t)(v')^p + t(v')^{2p})$$

when $0 \le t \le p-1$ and it is equal to

$$(1+v')^p = (1+(v')^p) = (1+(1+t)(v')^p + t(v')^{2p})$$

when $t = p$. □

Now we will calculate the total Chern class of the bundle over $B\Sigma_t$ that pulls backs to the normal bundle.

Lemma 3.5. *The total Chern class of* ξ_{Σ_t} *is*

$$\mathrm{c}(\xi_{\Sigma_t}) = 1 + (v')^{p-1} + \text{ \textit{higher terms}}$$

Proof. By Lemma 3.4 we know that the total Chern class of ϖ_{Σ_t} is

$$\mathrm{c}(\varpi_{\Sigma_t}) = 1 - (v')^{p-1} + \text{ higher terms}$$

By the construction of ξ_{Σ_t}, we know that $\xi_{\Sigma_t} \oplus \varpi_{\Sigma_t}$ is a trivial bundle over $B\Sigma_t^{(4p-1)}$, and the result follows. □

For the rest of this section set $r = \frac{p-1}{2}$.

Lemma 3.6. *The first few Pontrjagin classes of the bundle ξ_{Σ_t} are as follows*

$$p_k(\xi_{\Sigma_t}) = \begin{cases} 1 & \text{if } \ k = 0, \\ 0 & \text{if } \ 0 < k < r, \\ (-1)^r 2 (v')^{p-1} & \text{if } \ k = r \ . \end{cases}$$

Proof. This is direct calculation given Lemma 3.5 and the fact that

$$p_k(\xi_{\Sigma_t}) = \mathrm{c}_k(\xi_{\Sigma_t})^2 - 2\,\mathrm{c}_{k-1}(\xi_{\Sigma_t})\,\mathrm{c}_{k+1}(\xi_{\Sigma_t}) + - \cdots \mp 2\,\mathrm{c}_1(\xi_{\Sigma_t})\,\mathrm{c}_{2k-1}(\xi_{\Sigma_t}) \pm 2\,\mathrm{c}_{2k}(\xi_{\Sigma_t}) \ . \quad \square$$

The main result of this section is the following:

Lemma 3.7. $q_1(\xi_{\Sigma_t}) = v^{p-1} \in H^{2(p-1)}(B\Sigma_t; \mathbf{Z}/p)$

Proof. Let $\{K_n\}$ be the multiplicative sequence belonging to the polynomial $f(t) = 1 + t^r$. A result of Wu shows (see Theorem 19.7 in [30]) that

$$q_1(\xi_{\Sigma_t}) = K_r(p_1(\xi_{\Sigma_t}), \ldots, p_r(\xi_{\Sigma_t})) \bmod p \ .$$

By Lemma 3.6 we know that $p_1(\xi_{\Sigma_t}), \ldots, p_{r-1}(\xi_{\Sigma_t})$ are all zero, hence we are only interested in the coefficient of x_r in the polynomial $K_r(x_1, \ldots, x_r)$. By Problem 19-B in [30] this coefficient is equal to $s_r(0, 0, \ldots, 0, 1) = (-1)^{r+1} r$ (see [30, p. 188]) Hence we have

$$q_1(\xi_{\Sigma_t}) = (-1)^{r+1} r \bar{p}_r(\xi_{\Sigma_t}) = (-1)^{r+1} r (-1)^r 2 v^{p-1} = (-1)(p-1) v^{p-1} = v^{p-1}$$

where $\bar{p}_r(\xi_{\Sigma_t})$ denotes the mod p reduction of $p_r(\xi_{\Sigma_t})$. □

4. SMOOTH MODELS M_t AND N_t

Here we construct free smooth actions of the subgroups Σ_t and D_t on $S^{2p-1} \times S^{2p-1}$ and S^{4p-3} respectively, with the right bundle data. These provide models for covering spaces of the actions we are trying to construct.

§**4A**. **Construction of the examples M_t and N_t.** Given an m–dimensional representation $\Phi\colon G \to U(m)$ of a group G, we have an induced G–action on $\mathbb{C}^m$, and the space $S(\Phi) = S^{2m-1}$ will be the G–equivariant unit sphere in $\mathbb{C}^m$. We now construct two main examples.

(1) For $G = \Sigma_t$ and $t \in \{0, \ldots, p\}$, define

$$M_t = (S(\Psi_{\Sigma_t}) \times S((\Phi_{t,\Sigma_t})^{\oplus p})/\Sigma_t = (S^{2p-1} \times S^{2p-1})/\Sigma_t$$

(2) For $G = D_t$ and $t \in \{0, \dots, p\}$, define
$$N_t = S(\Phi_{t,D_t} \oplus \Phi^2_{t,D_t} \oplus \cdots \oplus \Phi^{p-1}_{t,D_t} \oplus (\Phi_{t,D_t})^{\oplus p})/D_t = S^{4p-3}/D_t$$
where, for a 1-dimensional representation $\mathit{\Phi}$, we set $\mathit{\Phi}^k$ to be the k^{th} power of $\mathit{\Phi}$ induced by the multiplication in S^1. In other words, if $\mathit{\Phi}(v) = \lambda v$ then we set $\mathit{\Phi}^k(v) = \lambda^k v$.

§4B. **The $(2p-1)$-type of M_t.** Let $X^{[n]}$ denote the n-type of a space X.

Lemma 4.1. *$M_t^{[2p-2]} \simeq B\Sigma_t$ and the composition $M_t \xrightarrow{i_{2p-2}} M_t^{[2p-2]} \xrightarrow{\simeq} B\Sigma_t$ is homotopy equivalent to the classifying map $c_t\colon M_t \to B\Sigma_t$.*

Proof. This follows as $S(\Psi_\Sigma, (\Phi_{t,\Sigma_t})^{\oplus p}) = S^{2p-1} \times S^{2p-1}$ is $(2p-2)$–connected and the action of Σ_t on $S(\Psi_{\Sigma_t}, (\Phi_{t,\Sigma_t})^{\oplus p})$ is free. ☐

Lemma 4.2. $M_t^{[2p-1]} \simeq B_{\Sigma_t}$.

Proof. Let $\mathrm{c}_p(\mathit{\Phi})$ denote the p^{th} Chern class of a representation $\mathit{\Phi}$ then
$$k_{2p-1}(M_t) = \mathrm{c}_p(\Psi_{\Sigma_t}) \oplus \mathrm{c}_p((\Phi_{t,\Sigma_t})^{\oplus p}) = \mathrm{Res}^\Gamma_{\Sigma_t}(\zeta) \oplus \mathrm{Res}^\Gamma_{\Sigma_t}(\alpha^p - \alpha^{p-1}\beta + \beta^p) = k_{\Sigma_t}$$
Hence the results follows by Lemma 4.1. ☐

§4C. **The tangent bundle of M_t.**

Lemma 4.3. *The tangent bundle of M_t is stably equivalent to the pull-back of $\tau_{\Sigma_t}\colon B_{\Sigma_t} \to BSO$ (see* Definition 3.1*).*

Proof. The tangent bundle $T(M_t)$ of M_t clearly fits into the following pull-back diagram
$$\begin{tikzcd}
T(M_t) \oplus \varepsilon \arrow[r] \arrow[d] & E\Sigma_t \times_{\Sigma_t} (\mathbb{C}^p \times \mathbb{C}^p) \arrow[d, "\pi"] \\
M_t \arrow[r, "c_t"] & B\Sigma_t
\end{tikzcd}$$
where ε is a trivial bundle over M_t and the action of Σ_t on $\mathbb{C}^p \times \mathbb{C}^p$ is given by Ψ_{Σ_t} and $(\Phi_{t,\Sigma_t})^{\oplus p}$ respectively. Hence we have $c_t^*([\pi]) = c_t^*([\psi_{\Sigma_t}] + p[\phi_{t,\Sigma_t}])$ in complex K-theory $\widetilde{K}(M_t)$. However, M_t fits into the following pull-back diagram:
$$\begin{tikzcd}
M_t \arrow[r] \arrow[d] & ED_t \times_{D_t} \mathbf{CP}^{p-1} \arrow[d] \\
L_t \arrow[r] & BD_t
\end{tikzcd}$$
where $L_t = S(\Phi^{\oplus p}_{t,D_t})/D_t$ and the action of D_t on $\mathbf{CP}^{p-1}$ is induced by action of D_t on $\mathbb{C}^p$ given by Ψ_{D_t}. Hence $\widetilde{K}(M_t)$ is a $\widetilde{K}(L_t)$–module by Proposition 2.13 in Chapter IV in [22] and the exponent of $\widetilde{K}(L_t)$ is p (see Theorem 2 in [21]). Hence the exponent of $\widetilde{K}(M_t)$ is p and $c_t^*([\pi]) = c_t^*([\psi_{\Sigma_t}]) = c_t^*([\varpi_{\Sigma_t}])$. This means the tangent bundle of M_t is stably equivalent to the pull-back of the bundle ϖ_{Σ_t} over $B\Sigma_t$ by the classifying map. However, by Lemma 4.1 and Lemma 4.2, we know that the classifying map is homotopy equivalent

to the compositon $M_t \xrightarrow{i_{2p-1}} M_t^{[2p-1]} \xrightarrow{\simeq} B_{\Sigma_t} \xrightarrow{\pi_{\Sigma_t}} B\Sigma_t$. Hence, the tangent bundle of M_t is stably equivalent to the pull-back of τ_{Σ_t}. □

§4D. The tangent bundle of N_t.

Lemma 4.4. *The tangent bundle of N_t is stably equivalent to the pull-back of $\varpi_{D_t}\colon BD_t \to BSO$.*

Proof. The tangent bundle $T(N_t)$ of N_t clearly fits into the following pull-back diagram

$$\begin{tikzcd}
T(N_t) \oplus \varepsilon \arrow[r] \arrow[d] & ED_t \times_{D_t} \mathbb{C}^{2p-1} \arrow[d] \\
N_t \arrow[r] & BD_t
\end{tikzcd}$$

where ε is a trivial bundle over N_t and the action of D_t on $\mathbb{C}^{2p-1}$ is given by $\Phi_{t,D_t} \oplus \Phi^2_{t,D_t} \oplus \cdots \oplus \Phi^{p-1}_{t,D_t} \oplus (\Phi_{t,D_t})^{\oplus p}$, where $\mathit{\Phi}^k(g) = (\mathit{\Phi}(g))^k$ for a 1-dimensional representation $\mathit{\Phi}\colon G \to S^1$. Now it is easy to see that

$$\varpi_{D_t} = 1 \oplus \Phi_{t,D_t} \oplus \Phi^2_{t,D_t} \oplus \cdots \oplus \Phi^{p-1}_{t,D_t} \oplus (\Phi_{t,D_t})^{\oplus p} \ .$$

Hence it is clear that the tangent bundle of N_t is the stably equivalent to the pull-back of ϖ_{D_t}. □

5. The image of the fundamental class

In this section we define a subset $T_\Gamma \subset H_{4p-3}(B_\Gamma; \mathbf{Z})$, which will turn out to contain the images of all the possible fundamental classes for our actions. To show that this set is non-empty, and consists of primitive elements of infinite order, we need to compute some cohomology groups of B_G, $G \subset \Gamma$. To carry out these computations, we will use the cohomology Serre spectral sequence of the fibration $K \longrightarrow B_G \xrightarrow{\pi_G} BG$, where $K = K(\mathbf{Z} \oplus \mathbf{Z}, 2p-1)$.

§5A. Definition of γ_{S^1}, γ_{Σ_t} and T_Γ. First note that the universal cover of M_t is $\mathbf{CP}^{p-1} \times S^{2p-1}$, for $0 \le t \le p$, and the universal cover of B_{Σ_t} is B_{S^1}. Hence, we can assume that we have the following pull-back diagram where the map c does not depend on t.

$$\begin{tikzcd}
\mathbf{CP}^{p-1} \times S^{2p-1} \arrow[r, "c"] \arrow[d] & B_{S^1} \arrow[d] \\
M_t \arrow[r, "c_t"] & B_{\Sigma_t}
\end{tikzcd}$$

We define an element γ_{S^1} in $H_{4p-3}(B_{S^1}; \mathbf{Z})$ as the image of the fundamental class of $\mathbf{CP}^{p-1} \times S^{2p-1}$ under the map c defined in the above diagram. In other words

$$\gamma_{S^1} = c_*\big[\mathbf{CP}^{p-1} \times S^{2p-1}\big] \in H_{4p-3}(B_{S^1}; \mathbf{Z}) \ .$$

Similarly, we define $\gamma_{\Sigma_t} \in H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$, $0 \le t \le p$, as the image of the fundamental class of M_t. In other words

$$\gamma_{\Sigma_t} = (c_t)_*\big[M_t\big] \in H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$$

Definition 5.1. We define

$$T_\Gamma = \{\gamma \in H_{4p-3}(B_\Gamma; \mathbf{Z}) \mid \ p \cdot (\mathrm{tr}(\gamma) - \gamma_{S^1}) = 0\}$$

where $\mathrm{tr}\colon H_{4p-3}(B_\Gamma; \mathbf{Z}) \to H_{4p-3}(B_{S^1}; \mathbf{Z})$ denotes the transfer map.

One of our main tasks will be to show that this subset T_Γ is non-empty !

§5B. **The (co)homology of K.** Let A be an abelian group. We will write

$${}_{(p)}A := A/\langle \text{torsion prime to } p\rangle \ .$$

The map

$$P^1\colon H^i(K; \mathbf{Z}/p) \to H^{i+2(p-1)}(K; \mathbf{Z}/p)$$

is the first mod p Steenrod operation and the map

$$\delta\colon H^i(K; \mathbf{Z}/p) \to H^{i+1}(K; \mathbf{Z})$$

is the Bockstein homomorphism. The following lemma gives the cohomology groups of K in the range we need.

Lemma 5.2. *Denote $H^{2p-1}(K; \mathbf{Z}) = \langle z_1, z_2\rangle = \mathbf{Z} \oplus \mathbf{Z}$ and let $\bar{z}_1$ and $\bar{z}_2$ be the mod p reductions of z_1 and z_2 respectively. We have*

1. *$H^i(K; A)$ is a torsion group for $2p \le i \le 4p-3$.*
2. *${}_{(p)}H^i(K; A) = 0$ for $2p \le i \le 4p-4$.*
3. *${}_{(p)}H^{4p-3}(K; \mathbf{Z}) = 0$.*
4. *${}_{(p)}H^{4p-2}(K; \mathbf{Z}) = \langle z_1 \cup z_2, \delta(P^1(\bar{z}_1)), \delta(P^1(\bar{z}_2))\rangle = \mathbf{Z} \oplus \mathbf{Z}/p \oplus \mathbf{Z}/p$.*
5. *$H^{4p-1}(K; \mathbf{Z})$ has no p-torsion.*

Proof. See results of Cartan [12], [13]. ☐

We will also need some information about the homology of K.

Lemma 5.3. *We have*

1. *$H_{2p-1}(K; \mathbf{Z}) = \mathbf{Z} \oplus \mathbf{Z}$,*
2. *$H_{2p-1}(K, ; \mathbf{Z}/p) = \mathbf{Z}/p \oplus \mathbf{Z}/p$,*
3. *${}_{(p)}H_i(K; A)$ is 0 for $2p-1 < i < 4p-3$, and*
4. *${}_{(p)}H_{4p-3}(K; \mathbf{Z}) = \mathbf{Z}/p \oplus \mathbf{Z}/p$.*

Proof. See results of Cartan [12], [13]. ☐

§5C. **The (co)homology spectral sequences.** For any subgroup $G \subset \Gamma$, let R be a ring and

$$\{E^r_{n,m}(G, R), d^r\} \quad \text{and} \quad \{E_r^{n,m}(G, R), d_r\}$$

be the homology and the cohomology Serre spectral sequences (respectively) of the fibration

$$K \longrightarrow B_G \xrightarrow{\pi_G} BG$$

where $K = K(\mathbf{Z} \oplus \mathbf{Z}, 2p-1)$. The second page of these spectral sequence is given by:

$$E^2_{n,m}(G, R) = H_n(BG; H_m(K; R)) \quad \text{and} \quad E_2^{n,m}(G, R) = H^n(BG; H^m(K; R)),$$

and they converge to $H_*(B_G; R)$ and to $H^*(B_G; R)$ respectively. The filtration $F_*H_*(B_G; R)$ of $H_*(B_G; R)$ is given by

$$F_nH_{n+m}(B_G; R) = \operatorname{Im}\left\{H_{n+m}(B_G^{\{n\}}; R) \xrightarrow{(i_n)_*} H_{n+m}(B_G; R)\right\}$$

and the filtration $F^*H^*(B_G; R)$ of $H^*(B_G; R)$ is given by

$$F^nH^{n+m}(B_G; R) = \ker\left\{H^{n+m}(B_G; R) \xrightarrow{(i_n)^*} H^{n+m}(B_G^{\{n-1\}}; R)\right\}$$

respectively, where

$$B_G^{\{n\}} = \pi_G^{-1}(BG^{(n)}) \ .$$

When $R = \mathbf{Z}$, we will write $\{E^r_{n,m}(G), d_r\}$ and $\{E_r^{n,m}(G), d_r\}$ instead of $\{E^r_{n,m}(G, \mathbf{Z}), d_r\}$ and $\{E_r^{n,m}(G, \mathbf{Z}), d_r\}$ respectively. The cohomology groups for

$$E_2^{*,0}(G, R) = H^*(BG; R)$$

are given in Theorem 2.3, Theorem 2.4, and Remark 2.5, and the calculation of

$$E_2^{0,*}(G, R) = H^*(K; R)$$

is given in Lemma 5.2.

§5D. **Definition of z_G, z'_G, and Z_G.** Let G be a subgroup of Γ which contains S^1. We have

$$E_{2p}^{0,2p-1}(G) = H^{2p-1}(K; \mathbf{Z}) = \langle z_1, z_2\rangle$$

and we can assume that

$$d_{2p}(z_1) = \begin{cases} \zeta & \text{if } G = \Gamma, \\ (\tau')^p - (v')^{p-1}\tau' & \text{if } G = \Sigma_t, \\ \tau^p & \text{if } G = S^1 \end{cases} \quad \text{and} \quad d_{2p}(z_2) = \begin{cases} \alpha^p - \alpha^{p-1}\beta + \beta^p & \text{if } G = \Gamma, \\ (v')^p & \text{if } G = \Sigma_t, \\ 0 & \text{if } G = S^1 \end{cases}$$

where $d_{2p}(z_1)$ and $d_{2p}(z_2)$ are in $E_{2p}^{2p,0}(G) = H^{2p}(BG; \mathbf{Z})$. Hence there exists z_G in $H^{2p-1}(B_G; \mathbf{Z})$ such that

$$H^{2p-1}(B_G; \mathbf{Z}) = \langle z_G\rangle = \mathbf{Z} \quad \text{and} \quad i^*(z_G) = \begin{cases} pz_2 & \text{if } G = \Gamma, \\ pz_2 & \text{if } G = \Sigma_t, \\ z_2 & \text{if } G = S^1 \end{cases}$$

where $i\colon K \to B_G$ is the inclusion map. Moreover we have

$$H^{2p-2}(B_G; \mathbf{Z}) = H^{2p-2}(B_G; \mathbf{Z}) = \begin{cases} \langle \alpha^{p-1}, \alpha^{p-2}\beta, \ldots, \beta^{p-1}, \chi_{p-1}\rangle & \text{if } G = \Gamma, \\ \langle (v')^{p-1}, (v')^{p-2}\tau', \ldots, (\tau')^{p-1}\rangle & \text{if } G = \Sigma_t, \\ \langle \tau^{p-1}\rangle & \text{if } G = S^1 \end{cases}$$

Define z'_G in $H^{2p-2}(B_G, \mathbf{Z})$ as follows

$$z'_G = \begin{cases} (\pi_\Gamma)^*(\chi_{p-1}) & \text{if } G = \Gamma, \\ (\pi_{\Sigma_t})^*((\tau')^{p-1}) & \text{if } G = \Sigma_t, \text{ and} \\ (\pi_{S^1})^*(\tau^{p-1}) & \text{if } G = S^1 \end{cases}$$

where $\pi_G : B_G \to BG$. Note that z'_G is a primitive element and generates a $\mathbf{Z}$ component in $H^{2p-2}(B_G;\mathbf{Z})$, where

$$H^{2p-2}(B_G;\mathbf{Z}) = \begin{cases} (\mathbf{Z}/p)^{\oplus p} \oplus \mathbf{Z} & \text{if } G = \Gamma, \\ (\mathbf{Z}/p)^{\oplus p} \oplus \mathbf{Z} & \text{if } G = \Sigma_t, \\ \mathbf{Z} & \text{if } G = S^1 \end{cases}$$

Define the *cohomology fundamental class* Z_G as follows:

$$Z_G = z_G \cup z'_G \in H^{4p-3}(B_G, \mathbf{Z}) \ .$$

The reason for this definition will become clear after Lemma 5.4. In the spectral sequence for $H^*(B_G;\mathbf{Z})$, $G = \Sigma_t$, the cohomology fundamental class of the manifold M_t lies in the term $E_\infty^{2p-2,2p-1}(G)$. In the formula for this term we see the elements z_G and z'_G described above.

§5E. **Transfers and duality.** In this section we will see the duality between Z_{S^1}, Z_{Σ_t}, and Z_Γ and γ_{S^1}, γ_{Σ_t}, and elements in T_Γ respectively. We will consider the p-fold covering maps

$$B_{\Sigma_t} \to B_\Gamma \ , \quad B_{S^1} \to B_{\Sigma_t} \ , \quad B\Sigma_t \to B\Gamma \ , \text{ and } \quad BS^1 \to B\Sigma_t$$

We will write π^* and π_* to denote the natural maps induced in cohomology and homology respectively, and just write *tr* for the transfer maps both in cohomology and homology. We have

$$\pi^*(z'_{\Sigma_t}) = z'_{S^1} \quad \text{and} \quad tr(z_{S^1}) = z_{\Sigma_t}$$

and

$$\pi^*(z_\Gamma) = z_{\Sigma_t} \quad \text{and} \quad tr(z'_{\Sigma_t}) = z'_\Gamma$$

Hence we have

$$tr(Z_{S^1}) = Z_{\Sigma_t} \quad \text{and} \quad tr(Z_{\Sigma_t}) = Z_\Gamma$$

Note that $\mathbf{CP}^{p-1} \times S^{2p-1}$ is the universal covering of M_t and we have the following pull-back diagram.

$$\begin{tikzcd} \mathbf{CP}^{p-1} \times S^{2p-1} \arrow[r, "c"] \arrow[d] & B_{S^1} \arrow[d] \\ M_t \arrow[r, "c_t"] & B_{\Sigma_t} \end{tikzcd}$$

Hence we have

$$\mathrm{tr}(\gamma_{\Sigma_t}) = \gamma_{S^1} \ .$$

Considering the map c: $\mathbf{CP}^{p-1} \times S^{2p-1} \to B_{S^1}$ we have

$$c^*(Z_{S^1}) = A \times B$$

where A is the cohomology fundamental class of $\mathbf{CP}^{p-1}$ and B is the cohomology fundamental class of S^{2p-1}. This proves that Z_{S^1} is a primitive element in $H^{4p-3}(B_{S^1};\mathbf{Z})$. Moreover

$$\langle Z_{S^1}, \gamma_{S^1} \rangle = \langle Z_{S^1}, c_*([\mathbf{CP}^{p-1} \times S^{2p-1}]) \rangle = \langle A \times B, [\mathbf{CP}^{p-1}] \times [S^{2p-1}] \rangle = 1 \ .$$

Hence we have

$$\langle Z_{\Sigma_t}, \gamma_{\Sigma_t} \rangle = \langle \operatorname{tr}(Z_{S^1}), \gamma_{\Sigma_t} \rangle = \langle Z_{S^1}, \operatorname{tr}(\gamma_{\Sigma_t}) \rangle = 1$$

and Z_{Σ_t} is a primitive element in $H^{4p-3}(B_{\Sigma_t}; \mathbf{Z})$. Moreover γ_{Σ_t}, and γ_{S^1} are primitive elements in $H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$ and $H_{4p-3}(B_{S^1}; \mathbf{Z})$ respectively. Hence our main calculation is the following: Given $\gamma \in T_\Gamma$, we have

$$\langle Z_\Gamma, \gamma \rangle = \langle \operatorname{tr}(Z_{S^1}, \gamma \rangle = \langle Z_{S^1}, \operatorname{tr}(\gamma) \rangle = \langle Z_{S^1}, \gamma_{S^1} \rangle = 1$$

since $\operatorname{tr}(\gamma) - \gamma_{S^1}$ is a torsion element.

§5F. Some spectral sequence calculations.

Lemma 5.4. *For $G = S^1$, Σ_t, or Γ, the differential $d_{2p}\colon E_{2p}^{2p-2,2p-1}(G) \to E_{2p}^{4p-2,0}(G)$ is surjective and its kernel is given as follows:*

$$E_\infty^{2p-2,2p-1}(G) = \begin{cases} \langle pz_2 \cdot \chi_{p-1} \rangle & \text{if } G = \Gamma, \\ \langle pz_2 \cdot (\tau')^{p-1} \rangle & \text{if } G = \Sigma_t, \\ \langle z_2 \cdot \tau^{p-1} \rangle & \text{if } G = S^1 \end{cases}$$

Proof. We have $E_{2p}^{2p-2,2p-1}(S^1) = \langle z_1 \cdot \tau^{p-1}, z_2 \cdot \tau^{p-1} \rangle = \mathbf{Z}^{\oplus 2}$ and $E_{2p}^{4p-2,0}(S^1) = \langle \tau^{2p-1} \rangle = \mathbf{Z}$. So result follows for $G = S^1$ because $d_{2p}(z_1 \cdot \tau^{p-1}) = \tau^{2p-1}$ spans $E_{2p}^{4p-2,0}(S^1)$. We have

$$E_{2p}^{2p-2,2p-1}(\Sigma_t) = (\mathbf{Z}/p)^{\oplus 2p} \oplus \mathbf{Z}^{\oplus 2}$$

given by

$$\langle z_1, z_2 \rangle \cdot \langle (v')^{p-1}, (v')^{p-2}\tau', \ldots, (\tau')^{p-1} \rangle \ .$$

and we have

$$E_{2p}^{4p-2,0}(\Sigma_t) = (\mathbf{Z}/p)^{\oplus 2p+1} \oplus \mathbf{Z}$$

given by

$$\langle (v')^{2p-1}, (v')^{2p-2}\tau', \ldots, (\tau')^{2p-1} \rangle \ .$$

The map

$$d_{2p}\colon E_{2p}^{2p-2,2p-1}(\Sigma_t) \to E_{2p}^{4p-2,0}(\Sigma_t)$$

is surjective because the following list of images of d_{2p} will span $E_{2p}^{4p-2,0}(\Sigma_t)$ considered as above:

- $d_{2p}(z_2 \cdot (v')^s (\tau')^{p-1-s}) = (v')^{p+s}(\tau')^{p-1-s}$ for $0 \le s \le p-1$
- $d_{2p}(z_1 \cdot (v')^s (\tau')^{p-1-s}) = (v')^s(\tau')^{2p-1-s} + (v')^{p-1+s}(\tau')^{p-s}$ for $0 \le s \le p-1$

This means we have

$$E_{2p+1}^{2p-2,2p-1}(\Sigma_t) = \langle pz_2 \cdot (\tau')^{p-1} \rangle$$

and the results follows for $G = \Sigma_t$, since $\ker d_{2p} = \langle pz_2 \cdot (\tau')^{p-1} \rangle$. The proof for $G = \Gamma$ is left to reader as it will not be used in this paper. □

Lemma 5.5. *$H^{4p-3}(B_{S^1}; \mathbf{Z}) = \langle Z_{S^1} \rangle \oplus A$ where A is a torsion group with no p-torsion.*

Proof. By Lemma 5.4

$$F^{2p-1}H^{4p-3}(B_{S^1};\mathbf{Z})=0$$

and

$$E_\infty^{2p-2,2p-1}(S^1)=\langle z_2\cdot\tau^{p-1}\rangle\ .$$

It is clear that

$$Z_{S^1}\in F^{2p-2}H^{4p-3}(B_{S^1};\mathbf{Z})$$

and represents the following generator of the quotient

$$z_2\cdot\tau^{p-1}\in E_\infty^{2p-2,2p-1}(S^1)=F^{2p-2}H^{4p-3}(B_{S^1};\mathbf{Z})/F^{2p-1}H^{4p-3}(B_{S^1};\mathbf{Z})\ .$$

Hence $F^{2p-2}H^{4p-3}(B_{S^1};\mathbf{Z})=\langle Z_{S^1}\rangle=\mathbf{Z}$, and the fact Z_{S^1} is a primitive element tells us that

$$H^{4p-3}(B_{S^1};\mathbf{Z})=\langle Z_{S^1}\rangle\oplus A$$

where A is an abelian group. Now by Lemma 5.2, we see that $E_\infty^{i,4p-3-i}(S^1)$ is a torsion group with no p-torsion for $0\le i\le 2p-3$. Hence A is a torsion group with no p-torsion. ☐

Lemma 5.6. *For $G=S^1$ or Σ_t, we have ${}_{(p)}H^{4p-2}(B_G;\mathbf{Z})=\mathbf{Z}/p\oplus\mathbf{Z}/p$.*

Proof. Note that $H^{2p-1}(B_G;\mathbf{Z})=0$ hence by Lemma 5.4 and Lemma 5.2, one can see that $E_\infty^{r,4p-2-r}(G)$ has no p-torsion for $1\le r\le 4p-2$, and the p-torsion part of $E_\infty^{0,4p-2}(G)$ is $\mathbf{Z}/p\oplus\mathbf{Z}/p$. ☐

Lemma 5.7. *The p-torsion subgroup of $H_{4p-3}(B_{\Sigma_t};\mathbf{Z}))$ is contained in the image of the natural map $i_*\colon H_{4p-3}(K;\mathbf{Z})\to H_{4p-3}(B_{\Sigma_t};\mathbf{Z}))$.*

Proof. By duality and Lemma 5.4 we see that the torsion part of $H_{4p-3}(B_{\Sigma_t};\mathbf{Z}))$ is equal to $F_{2p-3}H_{4p-3}(B_{\Sigma_t};\mathbf{Z})$. Hence the results follows from Lemma 5.3. ☐

Lemma 5.8. *The natural map $i_*\colon H_{2p-1}(K;\mathbf{Z}/p)\to H_{2p-1}(B_{\Sigma_t};\mathbf{Z}/p)$ is zero.*

Proof. It is clear that $i^*\colon H^{2p-1}(B_{\Sigma_t};\mathbf{Z}/p)\to H^{2p-1}(K;\mathbf{Z}/p)$ is zero. Because both $\bar z_1$ and $\bar z_2$ in

$$E_2^{0,2p-1}(\Sigma_t,\mathbf{Z}/p)=H^{2p-1}(K;\mathbf{Z}/p)=\langle\bar z_1,\bar z_2\rangle=\mathbf{Z}/p\oplus\mathbf{Z}/p$$

trangresses to nonzero elements in $E_2^{2p,0}(\Sigma_t,\mathbf{Z}/p)$. ☐

§5G. The subset T_Γ is non-empty. Let $\mathrm{tr}\colon H_{4p-3}(B_\Gamma;\mathbf{Z})\to H_{4p-3}(B_{S^1};\mathbf{Z})$ denote the transfer map.

Lemma 5.9. *Let γ' in $H_{4p-3}(B_\Gamma;\mathbf{Z})$ be an element such that $\langle Z_{S^1},\mathrm{tr}(\gamma')\rangle=1$. Then there exists an integer $N_{\gamma'}$ such that $p(1-p^2N_{\gamma'})(\mathrm{tr}(\gamma')-\gamma_{S^1})=0$.*

Proof. First note that $\langle Z_{S^1},\mathrm{tr}(\gamma')-\gamma_{S^1}\rangle=0$ since $\langle Z_{S^1},\gamma_{S^1}\rangle=1$. Hence $\mathrm{tr}(\gamma')-\gamma_{S^1}$ is a torsion element, by the Universal Coefficient Theorem and Lemma 5.5. Hence it is enough to prove that the order of $p\cdot(\mathrm{tr}(\gamma')-\gamma_{S^1})$ is relatively prime to p. But this is clear as the p-torsion of part of $H_{4p-3}(B_{S^1};\mathbf{Z})$ is same as the p-torsion part of $H^{4p-2}(B_{S^1};\mathbf{Z})$, which is $\mathbf{Z}/p\oplus\mathbf{Z}/p$ by Lemma 5.5. ☐

Theorem 5.10. *The set T_Γ is not empty. Any $\gamma \in T_\Gamma$ is a primitive element of infinite order in $H_{4p-3}(B_\Gamma; \mathbf{Z})$.*

Proof. Let tr denote the (co)homology transfer associated to the covering map $B_{S^1} \xrightarrow{\pi} B_\Gamma$. We know that $\mathrm{tr}(Z_{S^1}) = Z_\Gamma$ and Z_Γ is a primitive element in $H^{4p-3}(B_\Gamma; \mathbf{Z})$. Hence by the Universal Coefficient Theorem there exists a primitive element γ' in $H_{4p-3}(B_\Gamma; \mathbf{Z})$ such that $\mathrm{tr}(\gamma')$ is a primitive element in $H_{4p-3}(B_{S^1}; \mathbf{Z})$ and

$$\langle Z_\Gamma, \gamma' \rangle = 1 \text{ and } \langle Z_{S^1}, \mathrm{tr}(\gamma') \rangle = 1 \ .$$

Take $N_{\gamma'}$ as in Lemma 5.9. Define

$$\gamma_\Gamma = \gamma' - N_{\gamma'}(\pi)_*(\mathrm{tr}(\gamma') - \gamma_{S^1}) \ .$$

Then γ_Γ is in T_Γ, because by Lemma 5.9 we have

$$p \cdot (\mathrm{tr}(\gamma_\Gamma) - \gamma_{S^1}) = p(\mathrm{tr}(\gamma' - N_{\gamma'}(\pi)_*(\mathrm{tr}(\gamma') - \gamma_{S^1})) - \gamma_{S^1}) = p(1 - p^2 N_{\gamma'})(\mathrm{tr}(\gamma') - \gamma_{S^1}) = 0$$

Now, take any γ in T_Γ. Suppose that $\gamma = r\gamma_1$, for some γ_1 in $H_{4p-3}(B_\Gamma; \mathbf{Z})$. Then we would have $p \cdot (\gamma_{S^1} - r \cdot \mathrm{tr}(\gamma_1)) = 0$. But $\langle Z_{S^1}, \gamma_{S^1} \rangle = 1$. Hence $r = \pm 1$. □

Proposition 5.11. *Let* $\mathrm{tr}\colon H_{4p-3}(B_\Gamma; \mathbf{Z}) \to H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$ *denote the transfer map. Then any γ in T_Γ satisfies the following equation $p(\mathrm{tr}(\gamma) - \gamma_{\Sigma_t}) = 0$.*

Proof. For any γ in T_Γ the image of $p \cdot (\mathrm{tr}(\gamma) - \gamma_{\Sigma_t})$ under the transfer map from $H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$ to $H_{4p-3}(B_{S^1}; \mathbf{Z})$ is 0, by definition of T_Γ and the fact that $\mathrm{tr}(\gamma_{\Sigma_t}) = \gamma_{S^1}$. Note that the kernel of the above transfer map is included in the p-torsion part of $H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$, as $B_{S^1} \to B_{\Sigma_t}$ is p-covering. By Lemma 5.5, the p-torsion part of $H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$ is $\mathbf{Z}/p \oplus \mathbf{Z}/p$ (which has exponent p). This proves the result. □

6. The construction of the bordism element

The next step in our argument is to study the bordism groups $\Omega_{4p-3}(B_\Gamma, \nu_\Gamma)$ of our normal $(2p-1)$-type. The main result of this section is Theorem 6.9, which proves that the image of the Hurewicz map

$$\Omega_{4p-3}(B_\Gamma, \nu_\Gamma) \to H_{4p-3}(B_\Gamma; \mathbf{Z})$$

contains the non-empty subset T_Γ (see Definition 5.1). The main difficulty in computing the bordism groups is dealing with p-torsion. We will primarily use the James spectral sequence (a variant of the Atiyah-Hirzebruch spectral sequence) associated to the fibration

$$* \longrightarrow B_\Gamma \longrightarrow B_\Gamma$$

with E^2-term

$$E^2_{n,m}(\nu_\Gamma) = H_n(B_\Gamma; \Omega^{fr}_m(*)),$$

where the coefficients $\Omega^{fr}_m(*) = \pi^S_m$ are the stable homotopy groups of spheres. In our range, the p-torsion in π^S_m occurs only for π^S_{2p-3} and π^S_{4p-5}, where the p-primary part is $\mathbf{Z}/p$ (see [33, p. 5] and Example 6.3). This means that, after localizing at p, there are only two possibly non-zero differentials with source at the $(4p-3, 0)$-position, namely d_{2p-2} and d_{4p-4}. To show that these differentials are in fact both zero, and to prove that all other differentials starting at the $(4p-3,0)$-position also vanish on T_Γ, we use two techniques:

(i) For the differentials d^r with $2 \leq r \leq 4p-5$, and d^{4p-3}, we compare the James spectral sequence for $\Omega_{4p-3}(B_{\Gamma,}\nu_\Gamma)$ to the ones for $\Omega_{4p-3}(B_{\Sigma_t}, \nu_{\Sigma_t})$ via transfer, and use naturality.

(ii) For the differential d^{4p-4} we compare the James spectral sequence for $\Omega_{4p-3}(B_{\Gamma,}\nu_\Gamma)$ to the James spectral sequences for the fibrations $B\Sigma_t \to B\Gamma \to B(\Gamma/\Sigma_t)$, and use naturality again.

In carrying out the second step, we will need to use the Adams spectral sequence to prove that the natural map from the p–component of $\Omega^{fr}_{4p-5}(*)$ to $\Omega_{4p-5}(B\Sigma_t, \xi_{\Sigma_t})$ is injective (see Theorem 6.5).

§6A. **The James Spectral Sequence.** Let $\{E^r_{n,m}(\nu)\}$ denote the James spectral sequence (see [36]) associated to a vector bundle ν over a base space B and the fibration

$$* \longrightarrow B \longrightarrow B$$

and denote the differentials of this spectral sequence by d^r. We know that the second page is given by

$$E^2_{n,m}(\nu) = H_n(B, \Omega^{fr}_m(*))$$

and the spectral sequence converges to

$$E^\infty_{n,m}(\nu) = F_n\Omega_{n+m}(B,\nu)/F_{n-1}\Omega_{n+m}(B,\nu)$$

where $B^{(n)}$ stands for the n^{th} skeleton of B and

$$F_n\Omega_{n+m}(B,\nu) = \operatorname{Im}(\Omega_{n+m}(B^{(n)}, \nu|_{B^{(n)}}) \to \Omega_{n+m}(B,\nu))$$

For $0 \leq t \leq p$, let

$$\operatorname{tr}_t \colon E^r_{n,m}(\nu_\Gamma) \to E^r_{n,m}(\nu_{\Sigma_t})$$

denote the transfer map.

§6B. **Calculation of d^r when $2 \leq r \leq 4p-5$.** Here we employ our first technique. We first need some information about the James spectral sequences for $\Omega_{4p-3}(B_{\Sigma_t,}\nu_{\Sigma_t})$.

Lemma 6.1. *For $2 \leq r \leq 4p-5$, the differential*

$$d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t})$$

is zero on $\operatorname{tr}_t(T_\Gamma)$, where T_Γ is considered as subgroup of $E^r_{4p-3,0}(\nu_\Gamma)$.

Proof. Assume $2 \leq r \leq 4p-5$. By Proposition 5.11, and the fact that $d^r(\gamma_{\Sigma_t}) = 0$, it is enough to show that $d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t})$ is zero on the p-torsion subgroup I_t of $E^r_{4p-3,0}(\nu_{\Sigma_t}) = H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})$. Let I be the p-torsion part of $E^r_{4p-3,0}(\nu_{\Sigma_t}|_K) = H_{4p-3}(K; \mathbf{Z})$. By Lemma 5.7, we have

$$I_t = \operatorname{Im}\{i_* \colon I \to H_{4p-3}(B_{\Sigma_t}; \mathbf{Z})\}\ .$$

We consider the cases (i) $2 \leq r \leq 2p-3$, (ii) $r = 2p-2$, and (iii) $2p-1 \leq r \leq 4p-5$ separately.

Case (i). The group $E^2_{4p-3-r,r-1}(\nu_{\Sigma_t}|_K)$ is p-torsion free for $2 \le r \le 2p-3$, it follows that the differential

$$d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}|_K) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t}|_K)$$

is zero on I. Hence the differential

$$d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t})$$

is zero on I_t, for $2 \le r \le 2p-3$.

Case (ii). Next we observe that the map $i_* \colon E^2_{2p-1,2p-3}(\nu_{\Sigma_t}|_K) \to E^2_{2p-1,2p-3}(\nu_{\Sigma_t})$, restricted to p-torsion, is just the natural map $i_* \colon H_{2p-1}(K; \mathbf{Z}/p) \to H_{2p-1}(B_{\Sigma_t}; \mathbf{Z}/p)$, which is zero by Lemma 5.8 Hence, the differential

$$d^{2p-2} \colon E^{2p-2}_{4p-3,0}(\nu_{\Sigma_t}) \to E^{2p-2}_{2p-1,2p-3}(\nu_{\Sigma_t})$$

is zero on I_t by naturality.

Case (iii). Finally, we note that I_t is all p–torsion, but for $2p-1 \le r \le 4p-5$, the group $E^2_{4p-3-r,r-1}(\nu_{\Sigma_t})$ is p–torsion free. Hence, for $2p-1 \le r \le 4p-5$, the differential

$$d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t})$$

is zero on I_t. □

Lemma 6.2. *For $2 \le r \le 4p-5$, the differential*

$$d^r \colon E^r_{4p-3,0}(\nu_\Gamma) \to E^r_{4p-3-r,r-1}(\nu_\Gamma)$$

is zero on T_Γ, where T_Γ is considered as a subgroup of $E^r_{4p-3,0}(\nu_\Gamma)$.

Proof. Assume $2 \le r \le 4p-5$. By Lemma 6.1 we know that

$$d^r \colon E^r_{4p-3,0}(\nu_{\Sigma_t}) \to E^r_{4p-3-r,r-1}(\nu_{\Sigma_t})$$

is zero for all $t \in \{0, 1, \dots, p\}$. Hence it is enough to show that

$$\bigoplus\nolimits_t \operatorname{tr}_t \colon E^r_{4p-3-r,r-1}(\Gamma) \to \bigoplus\nolimits_t E^r_{4p-3-r,r-1}(\Sigma_t)$$

is injective. The map tr_0 is clearly injective for $r \ne 2p-2$ because the p–component of $\Omega^*_{r-1}(*)$ is 0 and $\widetilde{B_{\Sigma_t}} \to \widetilde{B_\Gamma}$ is a p-covering map. Hence $\bigoplus_t \operatorname{tr}_t$ is injective for $r \ne 2p-2$. Now we know that $\widetilde{B_\Gamma}$ and $\widetilde{B_{\Sigma_t}}$ are $2p-2$ connected. Hence for $r = 2p-2$ the map tr_t is the usual transfer map $H_{2p-1}(B\Gamma; \mathbf{Z}/p) \to H_{2p-1}(B\Sigma_t; \mathbf{Z}/p)$. Hence, it is enough to show that the map

$$\bigoplus\nolimits_t \operatorname{tr}_t \colon H_{2p-1}(B\Gamma; \mathbf{Z}/p) \to \bigoplus\nolimits_t H_{2p-1}(B\Sigma_t; \mathbf{Z}/p)$$

is injective. Dually, this is equivalent to showing that

$$\bigoplus\nolimits_t \operatorname{tr}_t \colon \bigoplus\nolimits_t H^{2p-1}(B\Sigma_t; \mathbf{Z}/p) \to H^{2p-1}(B\Gamma; \mathbf{Z}/p)$$

is surjective. By Theorem 2.4, we know that

$$H^{2p-1}(B\Gamma; \mathbf{Z}/p) = \langle x^{p-1}y, x^{p-2}x'y, \dots, (x')^{p-1}y, (x')^{p-1}y' \rangle \ .$$

Under the Bockstein homomorphism, this can be identified with

$$V_{p+1} = \langle \alpha^p, \alpha^{p-1}\beta, \dots, \alpha\beta^{p-1}, \beta^p \rangle \subseteq H^{2p}(B\Gamma; \mathbf{Z})$$

and this identifcation is natural with respect to the action of the automorphisms $\mathrm{Aut}(\Gamma)$ acting through the induced map $\mathrm{Aut}(\Gamma) \to GL_2(p)$. In the statements of Theorem 1 and Theorem 2 of [25], Leary gives explicit formulas for the action of $\mathrm{Aut}(\Gamma)$ on the generators of the cohomology rings $H^*(B\Gamma; \mathbf{Z})$ and $H^*(B\Gamma; \mathbf{Z}/p)$. The point is that these cohomology generators are pulled back from the quotient group $\mathbf{Z}/p \times \mathbf{Z}/p$.

Hence the action of the automorphisms $\phi_A \in \mathrm{Aut}(\Gamma)$, defined in Remark 2.2 for all $A \in SL_2(p)$, gives the standard $SL_2(p)$-action on V_{p+1}. This module V_{p+1} is known to be an indecomposable $SL_2(p)$-module (see [15, 5.7]), and there is a short exact sequence

$$0 \to V_2 \to V_{p+1} \to V_{p-1} \to 0$$

of $SL_2(p)$-modules, where $V_2 = \langle \alpha^p, \beta^p \rangle$ has dimension 2 and V_{p-1} is irreducible.

Now the image of the map $\bigoplus_t \mathrm{tr}_t$ is invariant under all automorphisms of the group Γ. Hence it is enough to show that $\mathrm{Im}(\bigoplus_t \mathrm{tr}_t)$ projects non-trivially into V_{p-1}. However, the calculations of [25, p. 67] show that

$$\mathrm{tr}_p(\mathrm{Res}_{\Sigma_p}(y') \cdot \bar{\tau}^{p-1}) = y' \cdot \mathrm{tr}_p(\bar{\tau}^{p-1}) = y'(c_{p-1} + x^{p-1}) = -(x')^{p-1}y' + y'x^{p-1} \ .$$

After applying the Bockstein, this shows that the element $\beta^p - \beta\alpha^{p-1}$ is contained in the image of the transfer. Since this element is not contained in the submodule V_2, we are done. □

The remaining possibly non-zero differentials are d^{4p-4} and d^{4p-3}. The first one is handled by comparison with the fibrations

$$B\Sigma_t \to B\Gamma \to B(\Gamma/\Sigma_t)$$

but first we must show that the induced map on coefficients at the $(0, 4p-3)$-position is injective on the p-component. For this we use the Adams spectral sequence.

§6C. **The Adams spectral sequence.** Let X be a connective spectrum of finite type. We will write

$$X = \{X_n, i_n \,|\, n \geq 0\},$$

where each X_n is a space with a basepoint and $i_n \colon \Sigma X_n \to X_{n+1}$ is a basepoint preserving map. We will denote the Adams spectral sequnce for X as follows:

$$\{E_r^{n,m}(X), d_r\} \ .$$

The second page of this spectral sequence is given by

$$E_2^{n,m}(X) = \mathrm{Ext}_{\mathcal{A}_p}^{n,m}(H^*(X; \mathbf{Z}/p), \mathbf{Z}/p),$$

where $\mathcal{A}_p$ is the mod-p Steenrod algebra and $H^*(X; \mathbf{Z}/p)$ is considered as an $\mathcal{A}_p$-module. The differentials of this spectral sequence are as follows:

$$d_r \colon E_r^{n,m} \to E_r^{n+r, m+r-1}$$

for $r \geq 2$, and it converges to

$${}_{(p)}\pi_*^S(X) = \pi_*^S(X)/\langle \text{torsion prime to } p \rangle$$

with the filtration

$$\cdots \subseteq F^{2,*+2}(X) \subseteq F^{1,*+1}(X) \subseteq F^{0,*}(X) = {}_{(p)}\pi_*^S(X)$$

defined by:

$$F^{n,m}(X) = {}_{(p)}\operatorname{Im}\{\pi_m^S(X_n) \to \pi_{m-n}^S(X)\}$$

In other words,

$$E_\infty^{n,m}(X) = F^{n,m}(X)/F^{n+1,m+1}(X) \ .$$

Example 6.3. Take an $\mathcal{A}_p$–free resolution $F_*^{\mathbb{S}}$ of the sphere spectrum $\mathbb{S}$

$$\ldots \xrightarrow{\partial_3} F_2^{\mathbb{S}} \xrightarrow{\partial_2} F_1^{\mathbb{S}} \xrightarrow{\partial_1} F_0^{\mathbb{S}} \xrightarrow{\partial_0} H^*(\mathbb{S};\mathbf{Z}/p)$$

with the following properties:

- We have $\iota_0^{\mathbb{S}}$ in $F_0^{\mathbb{S}}$, such that $\partial_0(\iota_0^{\mathbb{S}})$ is a generator of $H^*(\mathbb{S};\mathbf{Z}/p) = \mathbf{Z}/p$.
- We have $\alpha_0^{\mathbb{S}}$ and $\alpha_{2p-3}^{\mathbb{S}}$ in $F_1^{\mathbb{S}}$, such that $\partial_1(\alpha_0^{\mathbb{S}}) = \beta(\iota_0^{\mathbb{S}})$ and $\partial_1(\alpha_{2p-3}^{\mathbb{S}}) = P^1(\iota_0^{\mathbb{S}})$ because $H^i(\mathbb{S};\mathbf{Z}/p) = 0$ for $i \geq 1$.
- We have $\beta_{4p-5}^{\mathbb{S}}$ in $F_2^{\mathbb{S}}$, where $\partial_2(\beta_{4p-5}^{\mathbb{S}}) = P^2(\alpha_0^{\mathbb{S}}) - P^1\beta(\alpha_{2p-3}^{\mathbb{S}}) + 2\beta P^1(\alpha_{2p-3}^{\mathbb{S}})$ because
$$P^2(\beta(\iota_0^{\mathbb{S}})) - P^1\beta(P^1(\iota_0^{\mathbb{S}})) + 2\beta P^1(P^1(\iota_0^{\mathbb{S}})) = 0 \ .$$

In the Adams spectral sequence that converges to the p–component of $\pi_*^S(\mathbb{S}) = \Omega_*^{fr}(*)$, the element $\beta_{4p-5}^{\mathbb{S}}$ must survive to the E^∞–term as there are no possible differentials. Hence we have the following:

(1) ${}_{(p)}\Omega_{2p-3}^{fr}(*) = \mathbf{Z}/p = \langle \alpha_{2p-3}^{\mathbb{S}} \rangle$
(2) ${}_{(p)}\Omega_{4p-5}^{fr}(*) = \mathbf{Z}/p = \langle \beta_{4p-5}^{\mathbb{S}} \rangle$

§6D. Cohomology of the Thom spectrum associated to ξ_G. Now take any $G \subseteq \Gamma$ and let $M\xi_G$ denote the Thom spectrum associated to the bundle ξ_G. Since the bundle ξ_G is fixed, for a given G, we will shorten the notation by writing $MG = M\xi_G$. As in the previous section, we will denote an $\mathcal{A}_p$–free resolution of $H^*(MG;\mathbf{Z}/p)$ as follows:

$$\ldots \xrightarrow{\partial_2} F_1^{MG} \xrightarrow{\partial_1} F_0^{MG} \xrightarrow{\partial_0} H^*(MG;\mathbf{Z}/p)$$

It is clear that, to understand these resolutions we must first understand the $\mathcal{A}_p$–module structure on the cohomology $H^*(MG;\mathbf{Z}/p)$ of these spectra. Let $U_G \in H^0(MG;\mathbf{Z}/p)$ denote the Thom class of the Thom spectrum MG. Then we can write

$$H^*(MG;\mathbf{Z}/p) = U_G \cdot H^*(BG;\mathbf{Z}/p) \ .$$

Moreover, for $G = S^1$ we will write

$$H^*(BS^1;\mathbf{Z}/p) = \mathbf{F}_p[\bar{\tau}],$$

and for $G = D_t$ we have

$$H^*(BD_t;\mathbf{Z}/p) = (\Lambda(u) \otimes \mathbf{F}_p[v]) \ .$$

Hence for $G = \Sigma_t$ we can consider

$$H^*(B\Sigma_t;\mathbf{Z}/p) = H^*(BD_t;\mathbf{Z}/p) \otimes H^*(BS^1;\mathbf{Z}/p) = (\Lambda(u) \otimes \mathbf{F}_p[v]) \otimes \mathbf{F}_p[\bar{\tau}]$$

Lemma 6.4. *For $G = S^1$, D_t, or Σ_t we have*

$$\beta(U_G) = 0, \quad P^1\beta(U_G) = 0, \quad \beta P^1(U_G) = 0, \quad \beta P^1\beta(U_G) = 0, \quad P^2(U_G) = 0$$

and

$$P^1(U_G) = \begin{cases} U_{\Sigma_t}v^{p-1} & \text{if } \ G = \Sigma_t, \\ U_{D_t}v^{p-1} & \text{if } \ G = D_t, \\ 0 & \text{if } \ G = S^1 \end{cases}$$

Proof. The Thom class U_G is the mod p reduction of an integral cohomology class, so $\beta(U_G) = 0$. By Lemma 3.7, $q_1(\xi_{\Sigma_t}) = v^{p-1}$. Since $P^1(U_{\Sigma_t}) = U_{\Sigma_t}v^{p-1}$, we obtain

$$P^1(U_{S^1}) = 0 \ \text{ and } \ P^1(U_{D_t}) = U_{D_t}v^{p-1}$$

by restriction to $H^*(BD_t; \mathbf{Z}/p)$ and $H^*(BS^1; \mathbf{Z}/p)$. For $G = D_t$ or Σ_t we have

$$\beta P^1(U_G) = \beta(U_G v^{p-1}) = \beta(U)v^{p-1} + U\beta(v^{p-1}) = 0 + 0 = 0$$

and it is clear that $\beta P^1(U_{S^1}) = 0$. By the Adem relations we have $P^2(U_G) = 2P^1P^1(U_G)$. Hence for $G = D_t$ or Σ_t we have

$$P^2(U_G) = 2P^1(U_G v^{p-1}) = 2(P^1(U_G)v^{p-1}) + U_G P^1(v^{p-1})) = 2(U_G v^{2p-2} - U_G v^{2p-2}) = 0$$

and it is clear that $P^2(U_{S^1}) = 0$. □

§6E. Calculation of d^{4p-4}. The inclusion of a point induces a natural map from $\Omega^{fr}_{4p-5}(*)$ to $\Omega_{4p-5}(B\Sigma_t, \xi_{\Sigma_t})$ for each of the subgroups Σ_t, $0 \le t \le p$.

Theorem 6.5. *The natural map $\Omega^{fr}_{4p-5}(*) \to \Omega_{4p-5}(B\Sigma_t, \xi_{\Sigma_t})$ is injective on the p–component.*

Proof. The generator of ${}_{(p)}\Omega^{fr}_{4p-5}(*)$ is represented by the class $\beta^{\mathbb{S}}_{4p-5}$ defined above. We will show that this element maps non-trivially in the Adams spectral sequence. Denote the elements of $H^*(M\Sigma_t; \mathbf{Z}/p)$, $H^*(MS^1; \mathbf{Z}/p)$, and $H^*(MD_t; \mathbf{Z}/p)$ as in Section **§6D**. It is straightforward to check the following:

- We have $\iota^{M\Sigma_t}_0$ and $\iota^{M\Sigma_t}_{2p-3}$ in $F^{M\Sigma_t}_0$ such that

$$\partial_0(\iota^{M\Sigma_t}_0) = U_{\Sigma_t} \quad \text{and} \quad \partial_0(\iota^{M\Sigma_t}_{2p-3}) = U_{\Sigma_t}uv^{(p-2)} \ .$$

- We have $\alpha^{M\Sigma_t}_0$ and $\alpha^{M\Sigma_t}_{2p-3}$ in $F^{M\Sigma_t}_1$ such that

$$\partial_1(\alpha^{M\Sigma_t}_0) = \beta(\iota^{M\Sigma_t}_0) \ \text{ and } \ \partial_1(\alpha^{M\Sigma_t}_{2p-3}) = P^1(\iota^{M\Sigma_t}_0) - \beta(\iota^{M\Sigma_t}_{2p-3})$$

because $\beta(U_{\Sigma_t}) = 0$ and $P^1(U_{\Sigma_t}) - \beta(Uuv^{(p-2)}) = 0$.

- We also have $\alpha^{M\Sigma_t}_{4p-5}$ in $F^{M\Sigma_t}_1$ such that

$$\partial_1(\alpha^{M\Sigma_t}_{4p-5}) = P^1\beta(\iota^{M\Sigma_t}_{2p-3})$$

because $P^1(\beta(U_{\Sigma_t}uv^{(p-2)})) = 0$.

- We have $\beta^{M\Sigma_t}_{4p-5}$ in $F^{M\Sigma_t}_2$ such that

$$\partial_2(\beta^{M\Sigma_t}_{4p-5}) = P^2(\alpha^{M\Sigma_t}_0) - P^1\beta(\alpha^{M\Sigma_t}_{2p-3}) + 2\beta P^1(\alpha^{M\Sigma_t}_{2p-3}) + 2\beta(\alpha^{M\Sigma_t}_{4p-5})$$

because

$$P^2(\beta(\iota_0^{M\Sigma_t}))-P^1\beta(P^1(\iota_0^{M\Sigma_t})-\beta(\iota_{2p-3}^{M\Sigma_t}))+2\beta P^1(P^1(\iota_0^{M\Sigma_t})-\beta(\iota_{2p-3}^{M\Sigma_t}))+2\beta P^1\beta(\iota_{2p-3}^{M\Sigma_t})=0\ .$$

Now we define a part of the chain map $F_*^{M\Sigma_t} \to F_*^{\mathbb{S}}$. We send

$$\iota_0^{M\Sigma_t} \mapsto \iota_0^{\mathbb{S}} \ \text{ and } \ \iota_{2p-3}^{M\Sigma_t} \mapsto 0\ .$$

Since $\beta(\iota_0^{M\Sigma_t}) \mapsto \beta(\iota_0^{\mathbb{S}})$ and $P^1(\iota_0^{M\Sigma_t}) - \beta(\iota_{2p-3}^{M\Sigma_t}) \mapsto P^1(\iota_0^{\mathbb{S}})$ we must have

$$\alpha_{2p-3}^{M\Sigma_t} \mapsto \alpha_{2p-3}^{\mathbb{S}} \ \text{ and } \ \alpha_{4p-5}^{M\Sigma_t} \mapsto 0\ .$$

Finally, we can send

$$\beta_{4p-5}^{M\Sigma_t} \mapsto \beta_{4p-5}^{\mathbb{S}}$$

and this definition proves the Theorem, as there are no differentials in this range. □

Remark 6.6. A similar technique can be used to prove that the natural map $\Omega_{10}^{fr}(*) \to \Omega_{10}(BS^1, \xi_{S^1})$ is injective on the 3-component. One constructs a chain map $F_*^{\mathbb{S}} \to F_*^{MS^1}$ in degrees ≤ 11, whose composite with the chain map induced by the natural map $H^*(MS^1; \mathbf{Z}/p) \to H^*(\mathbb{S}; \mathbf{Z}/p)$ is chain homotopic to the identity. The element $\beta_{10}^{\mathbb{S}}$ generating the 3-component of π_{10}^S arises from $P^2(\alpha_3^{\mathbb{S}})$ and the Adem relation $P^2P^1\iota_0^{\mathbb{S}} = 0$.

Lemma 6.7. *$d^{4p-4}\colon E_{4p-3,0}^{4p-4}(\nu_\Gamma) \to E_{1,4p-5}^{4p-4}(\nu_\Gamma)$ is zero.*

Proof. We consider the fibration

$$B\Sigma_t \longrightarrow B\Gamma \longrightarrow B(\Gamma/\Sigma_t)$$

for $0 \leq t \leq p$. This fibration induces a James spectral sequence $E_{*,*}^*(t)$ with differential denoted by d_t^* so that the second page is given by

$$E_{n,m}^2(t) = H_n(\Gamma/\Sigma_t, \Omega_m(B\Sigma_t, \xi_{\Sigma_t}))$$

and the spectral sequence converges to $\Omega_*(B\Gamma, \xi_\Gamma)$. Moreover, we have a natural map $E_{*,*}^{4p-4}(\nu_\Gamma) \to E_{*,*}^{4p-4}(t)$ due to the following map of fibrations.

$$\begin{tikzcd}
* \arrow[r] \arrow[d] & B_\Gamma \arrow[r] \arrow[d] & B_\Gamma \arrow[d] \\
B\Sigma_t \arrow[r] & B\Gamma \arrow[r] & B(\Gamma/\Sigma_t)
\end{tikzcd}$$

Theorem 6.5 (applied for $t = 0$ and $t = p$), and the detection of $H_1(\mathbf{Z}/p \times \mathbf{Z}/p; \mathbf{Z}/p)$ by cyclic quotients, shows that the following sum of two of these natural maps is injective

$$E_{1,4p-5}^{4p-4}(\nu_\Gamma) \to E_{1,4p-5}^{4p-4}(0) \oplus E_{1,4p-5}^{4p-4}(p)\ .$$

However, the differential $d_t^{4p-4}\colon E_{4p-3,0}^{4p-4}(t) \to E_{1,4p-5}^{4p-4}(t)$ is zero for both $t = 0$ and $t = p$, since the element $N_{p-t} \to BD_{p-t} \to B\Gamma$, for $t = 0, p$ (defined in Section **§4A**) is non-zero in $\Omega_{4p-3}(B\Gamma, \xi_\Gamma)$. This is because $[N_{p-t}] \in H_{4p-3}(BD_{p-t}; \mathbf{Z})$ is non-zero, and the inclusion $D_{p-t} \subset \Gamma$ is split on homology by projection to $\Gamma/\Sigma_t \cong D_{p-t}$. □

§6F. Calculation of d^{4p-3}. The last differential doesn't involve p-torsion in the target, and can be handled by one more transfer argument.

Lemma 6.8. *$d^{4p-3}\colon E^{4p-3}_{4p-3,0}(\nu_\Gamma) \to E^{4p-4}_{0,4p-4}(\nu_\Gamma)$ is zero on T_Γ where T_Γ is considered as a subgroup of $E^{4p-3}_{4p-3,0}(\nu_\Gamma)$.*

Proof. By Lemma 6.7 and the transfer map tr_t we see that the differential

$$d^{4p-4}\colon E^{4p-4}_{4p-3,0}(\nu_{\Sigma_t}) \to E^{4p-4}_{1,4p-5}(\nu_{\Sigma_t})$$

is zero on $\mathrm{tr}_t(T_\Gamma)$ where T_Γ is considered as subgroup of $E^{4p-4}_{4p-3,0}(\nu_\Gamma)$. Now the differential

$$d^{4p-3}\colon E^{4p-3}_{4p-3,0}(\nu_{\Sigma_t}) \to E^{4p-4}_{0,4p-4}(\nu_{\Sigma_t})$$

has to be zero on γ_{Σ_t} and on the p-torsion group $\mathrm{Im}\{H_{4p-3}(K) \to H_{4p-3}(B_{\Sigma_t};\mathbf{Z})\}$, by Proposition 5.11 and the fact that $E^{4p-4}_{0,4p-4}(\nu_{\Sigma_t})$ is p-torsion free: this term is a quotient of $H_0(B_{\Sigma_t};\Omega^{fr}_{4p-4}(*)) \cong \pi^S_{4p-4}$, which has no p-torsion. Hence the result follows. □

We have now proved the main result of this section.

Theorem 6.9. *The subset $T_\Gamma \neq \emptyset$ is contained in the image of the Hurewicz map $\Omega_{4p-3}(B_\Gamma,\nu_\Gamma) \to H_{4p-3}(B_\Gamma;\mathbf{Z})$.*

Proof. Lemma 6.2, Lemma 6.7 and Lemma 6.8 shows that all the differentials going out of $E^r_{4p-3,0}(\nu_\Gamma)$ in the James spectral sequence for ν_Γ are zero on T_Γ and the result follows. □

7. SURGERY ON THE BORDISM ELEMENT

In this section we fix an odd prime p, the integer $n = 2p-1$, and assume that G is a finite subgroup of Γ that maps surjectively onto the quotient Q_p of Γ by S^1. We have now completed the first two steps in the proof of Theorem A. We have shown that there is a non-empty subset T_Γ consisting of primitive elements of infinite order in $H_{2n-1}(B_\Gamma)$, and that this subset is contained in the the image of the Hurewicz map $\Omega_{2n-1}(B_\Gamma,\nu_\Gamma) \to H_{2n-1}(B_\Gamma;\mathbf{Z})$. We now define the subset

$$T_G = \{trf(\gamma) \in H_{2n}(B_G;\mathbf{Z}) \mid \gamma \in T_\Gamma\},$$

where $trf\colon H_{2n-1}(B_\Gamma;\mathbf{Z}) \to H_{2n}(B_G;\mathbf{Z})$ denotes the S^1-bundle transfer induced by the fibration $S^1 \to B_G \to B_\Gamma$. Now fix

$$\gamma_G \in T_G\ .$$

By definition $\gamma_G = trf(\gamma)$, for some $\gamma \in T_\Gamma$, so we can pull back the S^1-bundle over a manifold (provided by Theorem 6.9) whose fundamental class represents γ under the bordism Hurewicz map. Hence we have a bordism class

$$[M^{2n}, f] \in \Omega_{2n}(B_G,\nu_G) \ \text{ such that } \quad \gamma_G = f_*[M]\ .$$

Surgery will be used to improve the manifold M within its bordism class. Our first remark is that we may assume f is an n-equivalence (see [23, Cor. 1, p. 719]). In particular, $\pi_1(M) = G$, and $\pi_i(M) = 0$ for $2 \leq i < n$. In addition, the map $f_*\colon \pi_n(M) \to \pi_n(B_G)$ is surjective. We need to determine the structure of $\pi_n(M)$ as a $\mathbf{Z}G$-module. First by applying the construction of [7, p. 230] to the chain complex $C(\widetilde{B_G})$ we get two $\mathbf{Z}G$-chain complexes $C(\theta_1)$ and $C(\theta_2)$ as in [7] (see Cor. 4.5 and Remark 3, p. 231), and investigated further in [6], with the following properties:

(i) We have $\theta_1 = \operatorname{Res}^\Gamma_G(\zeta)$ and $\theta_2 = \operatorname{Res}^\Gamma_G(\alpha^p - \alpha^{p-1}\beta + \beta^p)$.

(ii) There is a $\mathbf{Z}G$-chain map

$$\psi_i \colon C_*(\widetilde{B_G}) \longrightarrow C(\theta_i), \quad \text{for } i = 1, 2.$$

(iii) $H_*(C(\theta_i); \mathbf{Z}) = H_*(S^n; \mathbf{Z})$ and $H^*(C(\theta_i); \mathbf{Z}) = H^*(S^n; \mathbf{Z})$, for $i = 1, 2$.

(iv) There exists $[C(\theta_i)]$ a generator of $H^n(C(\theta_i); \mathbf{Z}) = \mathbf{Z}$ such that

$$(\psi_i)^*([C(\theta_i)]) = z_i, \quad i = 1, 2$$

where $H^n(\widetilde{B_G}; \mathbf{Z}) \cong H^n(K; \mathbf{Z}) = \langle z_1, z_2 \rangle \cong \mathbf{Z} \oplus \mathbf{Z}$.

(v) All the modules in the chain complex

$$D_* = C(\theta_1) \otimes_{\mathbf{Z}} C(\theta_2)$$

are finitely-generated projective $\mathbf{Z}G$-modules.

We will compare this complex to the complex $C_*(S(\Psi_G)) \otimes_{\mathbf{Z}} C(\theta_2)$.

Lemma 7.1. *The modules $C_i(S(\Psi_G)) \otimes_{\mathbf{Z}} C_j(\theta_2)$ are finitely-generated, projective $\mathbf{Z}G$-modules.*

Proof. The module $C_j(\theta_2)$ is free for $j < n$ and $C_i(S(\Psi_G))$ is free for $i > 2$. For $i \leq 2$, $C_i(S(\Psi_G))$ is a direct sum of free modules and modules of the form $\mathbf{Z}[G/D_t]$ for some t. Hence it is enough to show that $\mathbf{Z}[G/D_t] \otimes_{\mathbf{Z}} C_n(\theta_2)$ is a projective module for each t. We will use the criterion of [11, VI,8.10]: projectivity follows from cohomological triviality. There is an exact sequence

$$0 \to L_{\theta_2} \to \Omega^{n+1}\mathbf{Z} \xrightarrow{\theta_2} \mathbf{Z} \to 0$$

and

$$0 \to L_{\theta_2} \to F_n \to C_n(\theta_2) \to 0$$

where F_n is a free $\mathbf{Z}G$-module. Therefore, it is enough to show that the cohomology groups

$$\widehat{H}^q(G; \mathbf{Z}[G/D_t] \otimes_{\mathbf{Z}} L_{\theta_2}) = 0, \quad \text{for large } q \in \mathbf{Z}\ .$$

But we have an isomorphism (using the complete Ext-theory)

$$\widetilde{\operatorname{Ext}}^q_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], L_{\theta_2}) \cong \widehat{H}^q(G; \mathbf{Z}[G/D_t] \otimes_{\mathbf{Z}} L_{\theta_2})$$

by [11, III,2.2]. Now the long exact sequence

$$\begin{tikzcd}
\widetilde{\operatorname{Ext}}^q_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], L_{\theta_2}) \arrow[r] & \widetilde{\operatorname{Ext}}^q_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], \Omega^{n+1}\mathbf{Z}) \arrow[r] \arrow[d, equal] & \widetilde{\operatorname{Ext}}^q_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], \mathbf{Z}) \\
 & \widetilde{\operatorname{Ext}}^{q-n-1}_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], \mathbf{Z}) \arrow[ur, "\cup\theta_2"'] &
\end{tikzcd}$$

combined with Shapiro's Lemma [11, III,6.2]

$$\widetilde{\operatorname{Ext}}^q_{\mathbf{Z}G}(\mathbf{Z}[G/D_t], \mathbf{Z}) = \widetilde{\operatorname{Ext}}^q_{\mathbf{Z}D_t}(\mathbf{Z}, \mathbf{Z}),$$

and the fact that $\operatorname{Res}^G_{D_t}(\theta_2) \in H^{n+1}(D_t;\mathbf{Z})$ is a generator, completes the proof. $\square$

Lemma 7.2. *D_* is chain homotopy equivalent to a finite free $\mathbf{Z}G$-chain complex*

Proof. As in [7] we have the following pushout diagram

$$\begin{tikzcd}
 & L_{\theta_1} \arrow[r, equal] \arrow[d] & L_{\theta_1} \arrow[d] & & & & \\
0 \arrow[r] & \Omega^{n+1}\mathbf{Z} \arrow[r] \arrow[d, "\theta_1"] & C_n(\widetilde{BG}) \arrow[r] \arrow[d] & C_{n-1}(\widetilde{BG}) \arrow[r] \arrow[d] & \cdots \arrow[r] & C_0(\widetilde{BG}) \arrow[r] \arrow[d] & \mathbf{Z} \arrow[r] \arrow[d, "id"] & 0 \\
0 \arrow[r] & \mathbf{Z} \arrow[r] & C_n(\widetilde{BG})/L_{\theta_1} \arrow[r] & C_{n-1}(\widetilde{BG}) \arrow[r] & \cdots \arrow[r] & C_0(\widetilde{BG}) \arrow[r] & \mathbf{Z} \arrow[r] & 0
\end{tikzcd}$$

where the lower row is the chain complex $C(\theta_1)$.

If one extends the identity on $\mathbf{Z}$'s on the right hand side of the diagram below to a chain map $C_*(\widetilde{BG}) \to C_*(S(\Psi_G))$ then the map on the left hand side $\Omega^{n+1}\mathbf{Z} \to \mathbf{Z}$ must also represent θ_1 in $H^{n+1}(BG)$.

$$\begin{tikzcd}
0 \arrow[r] & \Omega^{n+1}\mathbf{Z} \arrow[r] \arrow[d, "\theta_1"] & C_n(\widetilde{BG}) \arrow[r] \arrow[d] & C_{n-1}(\widetilde{BG}) \arrow[r] \arrow[d] & \cdots \arrow[r] & C_0(\widetilde{BG}) \arrow[r, "\varepsilon"] \arrow[d] & \mathbf{Z} \arrow[r] \arrow[d, "id"] & 0 \\
0 \arrow[r] & \mathbf{Z} \arrow[r] & C_n(S(\Psi_G)) \arrow[r] & C_{n-1}(S(\Psi_G)) \arrow[r] & \cdots \arrow[r] & C_0(S(\Psi_G)) \arrow[r, "\varepsilon"] & \mathbf{Z} \arrow[r] & 0
\end{tikzcd}$$

This is because the class $\zeta \in H^{2p}(B\Gamma;\mathbf{Z})$ is the unique cohomology class u in this dimension such that $\operatorname{Res}^\Gamma_{D_t}(u) = 0$ for $0 \leq t \leq p$, and $\operatorname{Res}^\Gamma_{S^1}(u) = \tau^p$. On the other hand, by construction each subgroup $D_t \cong \mathbf{Z}/p$ is an isotropy subgroup of the action on $S(\Psi_G)$. The fixed-point complex $C_*(S(\Psi_G)^{D_t})$ has the homology of an odd-dimensional sphere (of lower dimension). Therefore, after restriction to D_t we can lift the identity on $\mathbf{Z}$ using

$$C_0(\widetilde{BG}) = \mathbf{Z}G \xrightarrow{\varepsilon} \mathbf{Z} \subseteq C_0(S(\Psi_G)^{D_t}) \subset C_0(S(\Psi_G))$$

and this lifting extends to the zero map $\Omega^{n+1}\mathbf{Z} \to \mathbf{Z}$.

Notice that these diagrams provide the translation between equivalence classes of multiple extensions and cohomology classes, as described in [29, III, 6.4]. Since $C(\theta_1)$ and $C_*(S(\Psi_G))$ considered as n-fold extensions from $\mathbf{Z}$ to $\mathbf{Z}$ both represent the same cohomology class, there is a chain map

$$C(\theta_1) \to C_*(S(\Psi_G))$$

which induces a cohomology isomorphism. Hence by the Kunneth formula we have a cohomology isomorphism

$$C(\theta_1) \otimes_{\mathbf{Z}} C(\theta_2) \to C_*(S(\Psi_G)) \otimes_{\mathbf{Z}} C(\theta_2)$$

where all the modules in $C_*(S(\Psi_G)) \otimes C(\theta_2)$ are projective. Therefore we have a chain homotopy equivalence of finitely-generated projective $\mathbf{Z}G$-chain complexes

$$D_* \to C_*(S(\Psi_G)) \otimes_{\mathbf{Z}} C(\theta_2)\ .$$

However, in the chain complex $C_*(S(\Psi_G)) \otimes C(\theta_2)$, all possibly non-free-modules projective modules have the form $\mathrm{Ind}_{D_t}^G(C_i(S(\Psi_G)^{D_t})) \otimes_{\mathbf{Z}} C_n(\theta_2)$. Since the Euler characteristic $\chi(C_*(S(\Psi_G)^{D_t})) = 0$, the finiteness obstruction of $C_*(S(\Psi_G)) \otimes_{\mathbf{Z}} C(\theta_2)$ vanishes. ☐

Lemma 7.3. *Under the transfer* $\mathrm{tr}\colon H_{2n}(B_G;\mathbf{Z}) \to H_{2n}(\widetilde{B_G};\mathbf{Z})$*, the class* $\mathrm{tr}(\gamma_G)$ *corresponds to the standard hyperbolic form*

$$\mathbf{H}(\mathbf{Z}) = (\mathbf{Z} \oplus \mathbf{Z}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix})$$

on $\pi_n(B_G) = \mathbf{Z} \oplus \mathbf{Z}$ *under the identification* $H_{2n}(\widetilde{B_G};\mathbf{Z})/Tors \cong \Gamma(\mathbf{Z} \oplus \mathbf{Z})$ *with Whitehead's* Γ*-functor.*

Proof. Let $d = |G|/p^2$ denote the order of the centre of G. We have a commutative diagram

$$\begin{tikzcd}
S^1 \arrow[r, "d"] \arrow[d] & S^1 \arrow[r] \arrow[d] & B\mathbf{Z}/d \arrow[d] \\
\widetilde{B_G} \arrow[r] \arrow[d] & B_G \arrow[r] \arrow[d] & BG \arrow[d] \\
B_{S^1} \arrow[r] & B_\Gamma \arrow[r] & BQ
\end{tikzcd}$$

where $Q = \mathbf{Z}/p \times \mathbf{Z}/p$ and $\widetilde{B_G} = K(\mathbf{Z} \oplus \mathbf{Z}, n)$. This gives a commutative square

$$\begin{tikzcd}
H_{2n-1}(B_{S^1};\mathbf{Z}) \arrow[r, "trf"] & H_{2n}(\widetilde{B_G};\mathbf{Z}) \\
H_{2n-1}(B_\Gamma;\mathbf{Z}) \arrow[r, "trf"] \arrow[u, "tr"] & H_{2n}(B_G;\mathbf{Z}) \arrow[u, "tr"]
\end{tikzcd}$$

relating the S^1-bundle transfers and the universal covering transfers. The p-torsion subgroup of $H_{2n-1}(B_{S^1};\mathbf{Z})$ maps to zero under the S^1-bundle transfer, since $H_{2n}(K;\mathbf{Z})$ has no p-torsion, by Lemma 5.2. Therefore $tr(\gamma_G) = trf(\gamma_{S^1})$ is just the image of the fundamental class of $S^{2p-1} \times S^{2p-1}$ in $H_{2n}(\widetilde{B_G};\mathbf{Z}) = H_{2n}(K;\mathbf{Z})$. But $H_{2n}(K;\mathbf{Z})/Tors = \mathbf{Z}$ can be naturally identified with $\Gamma(\mathbf{Z} \oplus \mathbf{Z}) = \mathbf{Z}$, and under this identification the fundamental class of $S^{2p-1} \times S^{2p-1}$ corresponds to a generator, represented by the hyperbolic plane. ☐

Theorem 7.4. *The equivariant intersection form of* M *is in the following form*

$$(\pi_n(M), s_M) \cong \mathbf{H}(\mathbf{Z}) \perp (F, \lambda)$$

where (F, λ) *is a non-singular skew-hermitian form on a finitely-generated free* $\mathbf{Z}G$*-module.*

Proof. Let $\psi\colon C_*(\widetilde{M}) \to D_*$ be the following composition

$$C_*(\widetilde{M}) \to C_*(\widetilde{B_G}) \xrightarrow{\Delta} C_*(\widetilde{B_G}) \otimes C_*(\widetilde{B_G}) \xrightarrow{\psi_1 \otimes \psi_2} D_*$$

where Δ denotes the diagonal map. First note that $\psi_*\colon H_i(\widetilde{M}) \to H_i(D_*)$ is clearly surjective for $i < 2n$. Assume $\psi_*([\widetilde{M}]) = [D]$ then

$$1 = \langle z_1 \cup z_2, f_*([\widetilde{M}])\rangle = \langle [C(\theta_1)] \otimes [C(\theta_2)], [D]\rangle$$

by the Kunneth formula. Hence ψ_* is also is surjective for $i \geq 2n$. As the image of the fundamental class $[M]$ of maps to a generator of $H_{2n}(D_*)$. Hence the homology of the mapping cone $H_i(\psi)$ is zero for $i \neq n$, and $H_n(\psi) = P$ is a finitely generated projective $\mathbf{Z}G$-module. But P is stably free by Lemma 7.2. Hence $\pi_n(M) = \mathbf{Z} \oplus \mathbf{Z} \oplus P$ where P is a finitely generated . By stabilizing M with connected sums of copies of $S^{2p-1} \times S^{2p-1}$ we may assume that $\pi_n(M) = \mathbf{Z} \oplus \mathbf{Z} \oplus F$, where F is a finitely-generated free $\mathbf{Z}G$-module.

To show the splitting of the equivariant intersection form $(\pi_n(M), s_M)$ we consider the relation

$$\langle f^*(z_1) \cup f^*(z_2), [\widetilde{M}]\rangle = \langle z_1 \cup z_2, f_*[\widetilde{M}]\rangle$$

where z_1, z_2 are a symplectic basis for the form on $\pi_n(B_G)$. Therefore, by Lemma 7.3, the map $f^*\colon H^n(\widetilde{B_G}; \mathbf{Z}) \to H^n(\widetilde{M}; \mathbf{Z})$ gives an isometric embedding of the hyperbolic form $\mathbf{H}(\mathbf{Z})$ into s_M. Any such isometric embedding splits (see [19, Lemma 1.4]). Hence the result follows. □

We next observe that the equivariant intersection form $(\pi_n(M), s_M)$ has a quadratic refinement $\mu\colon \pi_n(M) \to \mathbf{Z}G/\{\nu + \bar{\nu}\}$, in the sense of [40, Theorem 5.2]. This follows because the universal covering $\widetilde{M}$ has stably trivial normal bundle (use the Browder-Livesay quadratic map [9, Lemma 4.5, 4.6] for the elements of order two in G). We therefore obtain an element (F, λ, μ) of the surgery obstruction group (see [40, p. 49] for the essential definitions). The Arf invariant of this form is the Arf invariant of the associated form $\epsilon_*(F, \lambda, \mu)$, where $\epsilon\colon \mathbf{Z}G \to \mathbf{Z}$ is the augmentation map. This invariant factors through

$$\Omega_{2n}(B_G, \nu_G) \to \Omega_{2n}(B_\Gamma, \nu_\Gamma) \xrightarrow{Arf} \mathbf{Z}/2$$

and hence is zero for our bordism element. We also need to check the discriminant of this form.

Lemma 7.5. *We obtain an element*

$$(F, \lambda, \mu) \in L'_{2n}(\mathbf{Z}G)$$

of the weakly-simple surgery obstruction group.

Proof. A non-singular, skew-hermitian quadratic form (F, λ, μ) represents an element in $L'_{2n}(\mathbf{Z}G)$ provided that its discriminant lies in $\ker(\mathrm{Wh}(\mathbf{Z}G) \to \mathrm{Wh}(\mathbf{Q}G))$. But the equivariant symmetric Poincaré chain complex $(C(M), \varphi_0)$ is chain equivalent, after tensoring with the rationals $\mathbf{Q}$, to the rational homology (see [32, §4]). Therefore the image of the discriminant of $(\pi_n(M) \otimes \mathbf{Q}, s_M)$ equals the image of the torsion of φ_0, which vanishes in Wh($\mathbf{Q}G$) because closed manifolds have simple Poincaré duality (see [40, Theorem 2.1]). ☐

The proof of Theorem A. Suppose that p is an odd prime. We now have a representative $[M, f]$ for our bordism element in $\Omega_{2n}(B_G, \nu_G)$ whose equivariant intersection form $(\pi_n(M), s_M)$ contains (F, λ, μ) as described above. However, an element in the surgery obstruction group $L'_{2n}(\mathbf{Z}G)$ is zero provided that its multisignature and ordinary Arf invariant both vanish This is a result of Bak and Wall for groups of odd order (see [39, Cor. 2.4.3]), and for odd order groups direct product with cyclic groups we apply [39, Theorem 2.4.2 and Cor. 3.3.3]. The multisignature invariant is trivial since M is a closed manifold [40, 13B]. The ordinary Arf invariant of the universal covering $\widetilde{M}$ vanishes since $2n = 4p - 2$ is not of the form $2^k - 2$ (a famous result of Browder [10]). We can now do surgery on the classifying map $f\colon M \to B_G$ respecting the bordism class in $\Omega_{2n}(B_G, \nu_G)$ to obtain a representative $[M, f]$ which has $\widetilde{M} = S^n \times S^n \# \Sigma$, where Σ is a homotopy $2n$-sphere. Since the p-primary component of Cok J starts in dimension $2p(p-1) - 2$ (see [33, p. 5]) we can eliminate this homotopy sphere by equivariant connected sum unless $p = 3$.

In case $p = 3$, we use Remark 6.6 to show that $\widetilde{M} = S^5 \times S^5$. The bordism element $[\widetilde{M}, \widetilde{f}] \in \Omega_{10}^{fr}(K)$ vanishes in $\Omega_{10}(B_{S^1}, \nu_{S^1})$ by the Gysin sequence in bordism. But the difference element $[\widetilde{M}, \widetilde{f}] - [S^5 \times S^5, i_5] \in \Omega_{10}^{fr}(*)$. Since $\Omega_{10}^{fr}(*)$ injects on the 3-component into $\Omega_{10}(BS^1, \xi_{S^1})$, it follows that the order of the difference element is not divisible by 3. Thus in all cases we can obtain $\widetilde{M} = S^n \times S^n$. This completes the proof of Theorem A. ☐

REFERENCES

[1] A. Adem, *Constructing and deconstructing group actions*, Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory, Contemp. Math., vol. 346, Amer. Math. Soc., Providence, RI, 2004, pp. 1–8.

[2] A. Adem, J. F. Davis, and Ö. Ünlü, *Fixity and free group actions on products of spheres*, Comment. Math. Helv. **79** (2004), 758–778.

[3] A. Adem and J. H. Smith, *Periodic complexes and group actions*, Ann. of Math. (2) **154** (2001), 407–435.

[4] K. Alzubaidy, *Free actions of p-groups* ($p > 3$) *on* $S^n \times S^n$, Glasgow Math. J. **23** (1982), 97–101.

[5] ———, *Free actions on* $(S^n)^k$, Mathematika **32** (1985), 49–54.

[6] D. J. Benson and J. F. Carlson, *Projective resolutions and Poincaré duality complexes*, Trans. Amer. Math. Soc. **342** (1994), 447–488.

[7] D. J. Benson and J. F. Carlson, *Complexity and multiple complexes*, Math. Z. **195** (1987), 221–238.

[8] N. Blackburn, *Generalizations of certain elementary theorems on p-groups*, Proc. London Math. Soc. (3) **11** (1961), 1–22.

[9] W. Browder and G. R. Livesay, *Fixed point free involutions on homotopy spheres*, Tôhoku Math. J. (2) **25** (1973), 69–87.

[10] W. Browder, *The Kervaire invariant of framed manifolds and its generalization*, Ann. of Math. (2) **90** (1969), 157–186.

[11] K. S. Brown, *Cohomology of groups*, Springer-Verlag, New York, 1994, Corrected reprint of the 1982 original.

[12] H. Cartan, *Sur les groupes d'Eilenberg-Mac Lane. II*, Proc. Nat. Acad. Sci. U. S. A. **40** (1954), 704–707.

[13] ———, *Séminaire Henri Cartan de l'Ecole Normale Supérieure, 1954/1955. Algèbres d'Eilenberg-MacLane et homotopie*, Secrétariat mathématique, 11 rue Pierre Curie, Paris, 1955.

[14] P. E. Conner, *On the action of a finite group on $S^n \times S^n$*, Ann. of Math. (2) **66** (1957), 586–588.

[15] D. J. Glover, *A study of certain modular representations*, J. Algebra **51** (1978), 425–475.

[16] I. Hambleton, *Some examples of free actions on products of spheres*, Topology **45** (2006), 735–749.

[17] I. Hambleton and O. Ünlü, *Examples of free actions on products of spheres*, Quart. J. Math. (to appear), arXiv:math.AT 0705.4081, 2008.

[18] A. Heller, *A note on spaces with operators*, Illinois J. Math. **3** (1959), 98–100.

[19] F. Hirzebruch, W. D. Neumann, and S. S. Koh, *Differentiable manifolds and quadratic forms*, Marcel Dekker Inc., New York, 1971, Appendix II by W. Scharlau, Lecture Notes in Pure and Applied Mathematics, Vol. 4.

[20] M. A. Jackson, Qd(p)*-free rank two finite groups act freely on a homotopy product of two spheres*, J. Pure Appl. Algebra **208** (2007), 821–831.

[21] T. Kambe, *The structure of K_Λ-rings of the lens space and their applications*, J. Math. Soc. Japan **18** (1966), 135–146.

[22] M. Karoubi, *K-theory*, Springer-Verlag, Berlin, 1978, An introduction, Grundlehren der Mathematischen Wissenschaften, Band 226.

[23] M. Kreck, *Surgery and duality*, Ann. of Math. (2) **149** (1999), 707–754.

[24] I. J. Leary, *The integral cohomology rings of some p-groups*, Math. Proc. Cambridge Philos. Soc. **110** (1991), 25–32.

[25] ———, *The mod-p cohomology rings of some p-groups*, Math. Proc. Cambridge Philos. Soc. **112** (1992), 63–75.

[26] I. J. Leary, *The cohomology of ceratin groups*, Ph.D. thesis, University of Cambridge, 1991, (http://www.maths.abdn.ac.uk∼bensondj/html/archive/leary.html).

[27] G. Lewis, *Free actions on $S^n \times S^n$*, Trans. Amer. Math. Soc. **132** (1968), 531–540.

[28] ———, *The integral cohomology rings of groups of order p^3*, Trans. Amer. Math. Soc. **132** (1968), 501–529.

[29] S. Mac Lane, *Homology*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1975 edition.

[30] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J., 1974, Annals of Mathematics Studies, No. 76.

[31] R. Oliver, *Free compact group actions on products of spheres*, Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), Lecture Notes in Math., vol. 763, Springer, Berlin, 1979, pp. 539–548.

[32] A. Ranicki, *The algebraic theory of surgery. I. Foundations*, Proc. London Math. Soc. (3) **40** (1980), 87–192.

[33] D. C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, AMS Chelsea Publishing, 2004.

[34] E. Stein, *Free actions on products of spheres*, Michigan Math. J. **26** (1979), 187–193.

[35] R. E. Stong, *Notes on cobordism theory*, Mathematical notes, Princeton University Press, Princeton, N.J., 1968.

[36] P. Teichner, *On the signature of four-manifolds with universal covering spin*, Math. Ann. **295** (1993), 745–759.

[37] C. B. Thomas, *Free actions by p-groups on products of spheres and Yagita's invariant po(G)*, Transformation groups (Osaka, 1987), Lecture Notes in Math., vol. 1375, Springer, Berlin, 1989, pp. 326–338.

[38] Ö. Ünlü, *Constructions of free group actions on products of spheres*, Ph.D . Thesis, University of Wisconsin, 2004.

[39] C. T. C. Wall, *Classification of Hermitian Forms. VI. Group rings*, Ann. of Math. (2) **103** (1976), 1–80.

[40] ———, *Surgery on compact manifolds*, second ed., American Mathematical Society, Providence, RI, 1999, Edited and with a foreword by A. A. Ranicki.

[41] N. Yagita, *On the dimension of spheres whose product admits a free action by a nonabelian group*, Quart. J. Math. Oxford Ser. (2) **36** (1985), 117–127.

Department of Mathematics & Statistics
McMaster University
Hamilton, ON L8S 4K1, Canada
E-mail address: `ian@math.mcmaster.ca`

Department of Mathematics & Statistics
McMaster University
Hamilton, ON L8S 4K1, Canada
E-mail address: `unluo@math.mcmaster.ca`